

GEOGRAPHICAL DISTRIBUTION.

THERE is given here a list of the Stations and other places from which the Challenger Expedition procured Scaphopods and Marine Gasteropods, so far, at least, as these have been sent to me. The names of these molluscs from each locality have been added in alphabetical order as simplest for reference, and all species first brought to light by the Expedition are here reckoned as new.

Station II. January 13, 1873. Lat. 38° 10' N., long. 9° 14' W. Off Setubal. 470 fathoms. Green mud.

- | | |
|--|--|
| <p><i>Bittium gemmatum</i>, n. sp., p. 547, pl. xxxix.
fig. 2.</p> <p><i>Columbella</i> (<i>Anachis</i>) <i>halicæsti</i>, Jeffr., p. 236.</p> <p><i>Dentalium capillosum</i>, Jeffr. var., p. 1, pl. i.
fig. 1.</p> <p>„ <i>circumcinctum</i>, n. sp., p. 7, pl. i. fig. 7.</p> | <p><i>Dentalium entalis</i>, Linne, var. <i>orthrum</i>, Wats., p. 6.</p> <p><i>Fusus</i> (<i>Neptunea</i>) <i>despectus</i> (Linne), p. 199.</p> <p>„ (<i>Siphonorbis</i>) <i>amblyterus</i>, n. sp., p. 205, pl. xvii.
fig. 8.</p> <p><i>Trochus</i> (<i>Margarita</i>) <i>rhysus</i>, n. sp., p. 83, pl. v. fig. 4.</p> |
|--|--|

February 1873. From the rocks. Santa Cruz. Tenerife.

- | | |
|---|---|
| <p><i>Litorina striata</i>, King, p. 575.</p> <p><i>Ranella</i> (<i>Lampas</i>) <i>thomæ</i>, d'Orb., p. 400.</p> | <p><i>Trochus</i> (<i>Trochocochlea</i>) <i>colubrinus</i>, Gould, p. 65.</p> |
|---|---|

Station VIIp. February 10, 1873. Lat. 28° 35' N., long. 16° 5' W. Off Tenerife. Canaries. 78 fathoms. Volcanic sand.

- | | |
|---|--|
| <p><i>Aclis mizon</i>, n. sp., p. 501, pl. xxxiv. fig. 1.</p> <p><i>Bittium lacteum</i> (Phil.), p. 539.</p> <p>„ <i>reticulatum</i> (Da Costa), p. 540.</p> <p><i>Cæcum elegantissimum</i>, Carp., p. 686.</p> <p>„ <i>pollicare</i>, Carp., p. 680.</p> <p>„ <i>vitreum</i>, Carp., p. 680.</p> <p><i>Columbella rustica</i> (Linne), p. 233.</p> <p><i>Cypræa spurca</i>, Linne, p. 425.</p> <p>„ (<i>Trivia</i>) <i>candidula</i>, Gask., p. 427.</p> <p><i>Eulima philippii</i>, Weink., p. 506.</p> <p><i>Fasciolaria armata</i>, A. Adams, p. 243, pl. xiii.
fig. 1.</p> <p><i>Marginella</i> (<i>Gibberula</i>) <i>guanacha</i>, d'Orb., p. 270.</p> <p>„ „ <i>miliaria</i> (Linne), p. 271.</p> <p><i>Natica variabilis</i>, Recl., p. 435.</p> <p>„ (<i>Mamma</i>) <i>porcellana</i>, d'Orb., p. 453.</p> <p><i>Odostomia turrata</i>, Hanley, p. 481.</p> <p><i>Phasianella pulla</i> (Linne), p. 173.</p> | <p><i>Philine aperta</i> (Linne), p. 671.</p> <p><i>Ranella</i> (<i>Aspa</i>) <i>marginata</i> (Gmel.), p. 403.</p> <p><i>Rissoa costulata</i>, Alder, p. 586.</p> <p>„ <i>similis</i>, Scacc., p. 587.</p> <p>„ (<i>Alvania</i>) <i>canariensis</i>, d'Orbigny, var. Manzoni,
p. 593.</p> <p>„ „ <i>costata</i> (J. Adams), p. 594.</p> <p>„ „ <i>leacocki</i>, Wats., p. 594.</p> <p>„ (<i>Oingula</i>) <i>glabrata</i> (v. Mühl.), p. 604.</p> <p>„ (<i>Onoba</i>) <i>striata</i> (J. Adams), p. 598.</p> <p>„ „ <i>watsoni</i>, Schwartz, p. 599.</p> <p>„ (<i>Setia</i>) <i>albugo</i>, Wats., p. 606.</p> <p>„ „ <i>callosa</i>, Manz., p. 606.</p> <p><i>Triforis perversa</i> (Linne), p. 560.</p> <p><i>Trochus</i> (<i>Zisiphinus</i>) <i>striatus</i>, Linne, p. 56.</p> <p><i>Turritella bicingulata</i>, Lam., p. 467.</p> <p><i>Utriculus mamillatus</i> (Phil.), p. 647.</p> <p>„ <i>tornatus</i>, n. sp., p. 651, pl. xlvi. fig. 10.</p> |
|---|--|