

- | | |
|--|---|
| <i>Neobuccinum vestitum</i> , v. Mart., p. 216. | <i>Struthiolaria mirabilis</i> , E. A. Sm., p. 414. |
| <i>Pleurotoma (Spirotropis) studeriana</i> , v. Mart.,
p. 322, pl. xxv. fig. 7. | <i>Trophon albolabratus</i> , E. A. Sm., p. 165. |
| „ <i>(Thesbia) translucida</i> , n. sp., p. 330,
pl. xxv. fig. 11. | <i>Trochus (Photinula) expansus</i> (Sow.), p. 69. |

Station 149C. January 19, 1874. Lat. 49° 32' S., long. 70° E. Balfour Bay, Royal Sound, Kerguelen. 60 fathoms. Volcanic mud.

- | | |
|--|--|
| <i>Actæon (Actæonina) edentula</i> , n. sp., p. 632,
pl. xlvi. fig. 6. | <i>Natica (Amauropsis) suturalis</i> , n. sp., p. 455, pl. xxvii.
fig. 4. |
| <i>Cancellaria (Admete)</i> , sp. | <i>Patella fuegiensis</i> , Reeve, p. 26. |
| <i>Fusus (Euthria) chloroticus</i> , v. Mart., p. 209,
pl. xviii. fig. 8. | <i>Puncturella noachina</i> (Linne), var. <i>princeps</i> , Migh.,
p. 42. |
| <i>Litorina setosa</i> , E. A. Sm., p. 575. | <i>Struthiolaria mirabilis</i> , E. A. Sm., p. 414. |
| <i>Natica fartilis</i> , n. sp., p. 446, pl. xxvii. fig. 10. | <i>Trophon albolabratus</i> , E. A. Sm., p. 165. |
| „ <i>(Lunatia) prasina</i> , n. sp., p. 449,
pl. xxvii. fig. 9. | <i>Trochus (Photinula) expansus</i> (Sow.), p. 69. |

January 19–20, 1874. Royal Sound, Kerguelen. Shore.

- | | |
|--|---|
| <i>Eatoniella caliginosa</i> (E. A. Sm.), p. 614. | <i>Rissoa (Setia) principis</i> , n. sp., p. 608, pl. xliv.
fig. 13. |
| <i>Hydrobia caliginosa</i> (Gould), p. 613. | „ „ <i>sinapi</i> , n. sp., p. 610, pl. xlvi. fig. 13. |
| <i>Patella fuegiensis</i> , Reeve, p. 26. | <i>Scissurella obliqua</i> , n. sp., p. 116, pl. viii. fig. 5. |
| <i>Rissoa (Setia) australis</i> , n. sp., p. 608, pl. xlvi.
fig. 1. | <i>Skenea subcanaliculata</i> , E. A. Sm., p. 624. |

January 1874. Kerguelen. Rocks.

- | | |
|--|---|
| <i>Neobuccinum eatoni</i> , E. A. Sm., p. 216. | <i>Patella kerguelensis</i> , E. A. Sm., p. 27. |
| <i>Patella fuegiensis</i> , Reeve, p. 26. | : |

Station 149D. January 20, 1874. Lat. 49° 28' S., long. 70° 13' E. Royal Sound, Kerguelen. 28 fathoms. Volcanic mud.

- | | |
|--|---|
| <i>Buccinum albozonatum</i> , n. sp., p. 212, pl. xiii. fig. 7. | <i>Pleurotoma (Spirotropis) studeriana</i> , v. Mart., p. 322,
pl. xxv. fig. 7. |
| <i>Cancellaria (Admete) carinata</i> , n. sp., p. 275,
pl. xviii. fig. 9. | „ <i>(Thesbia) translucida</i> , n. sp., p. 330,
pl. xxv. fig. 11. |
| <i>Fusus (Euthria) chloroticus</i> , v. Mart., p. 209,
pl. xviii. fig. 8. | „ „ <i>platamodes</i> , n. sp., p. 332,
pl. xxiv. fig. 3. |
| „ <i>(Neptunea) regulus</i> , n. sp., p. 204,
pl. xiii. fig. 7. | „ <i>(Typhlomangelia) fluctuosa</i> , n. sp.,
var. <i>cariosa</i> , p. 319, pl. xxv. fig. 4. |
| <i>Natica grisea</i> , v. Mart., p. 432, pl. xxviii. fig. 5. | „ sp. |
| „ <i>(Amauropsis) perscalpta</i> , v. Mart.,
p. 454, pl. xxviii. fig. 4. | <i>Struthiolaria mirabilis</i> , E. A. Sm., p. 414. |
| „ <i>(Lunatia) prasina</i> , n. sp., p. 449,
pl. xxvii. fig. 9. | <i>Trochus (Photinula) expansus</i> (Sow.), p. 69. |
| <i>Neobuccinum vestitum</i> , v. Mart., p. 216. | <i>Trophon albolabratus</i> , E. A. Sm., p. 165. |
| <i>Pleurotoma (Thesbia) corpulenta</i> , n. sp., p. 331,
pl. xxv. fig. 8. | „ <i>septus</i> , n. sp., p. 170, pl. x. fig. 11. |
| | <i>Turritella austrina</i> , n. sp., p. 470, pl. xxix. fig. 2. |
| | <i>Volutomitra fragillima</i> , n. sp., p. 263, pl. xiv. fig. 7. |