

PLATE XXV.

	PAGE
Fig. 1. <i>Pleurotoma (Surcula) gypsata</i> , n. sp. . . . .	297
<i>a.</i> The natural size. . . . .   <i>b.</i> The apex, enlarged.	
Fig. 2. <i>Pleurotoma (Surcula) plebeia</i> , n. sp. . . . .	292
<i>a.</i> The natural size. . . . .   <i>b.</i> The sculpture, enlarged.	
Fig. 3. <i>Pleurotoma (Typhlomangelia) paupera</i> , n. sp. . . . .	317
<i>a.</i> The natural size. . . . .   <i>b.</i> The sculpture, enlarged.	
Fig. 4. <i>Pleurotoma (Typhlomangelia) fluctuosa</i> , n. sp. . . . .	319
<i>a, b.</i> The shell, natural size. . . . .   <i>e.</i> The operculum, enlarged.	
<i>c.</i> The apex, enlarged. . . . .   <i>f.</i> Natural size of the same.	
<i>d.</i> The sculpture, enlarged. . . . .	
Fig. 5. <i>Pleurotoma (Surcula) trilix</i> , n. sp. . . . .	287
<i>a.</i> The shell, natural size. . . . .   <i>c.</i> The sculpture, enlarged.	
<i>b.</i> The apex, enlarged. . . . .	
Fig. 6. <i>Pleurotoma (Typhlomangelia) pyrrha</i> , n. sp. . . . .	316
<i>a, b.</i> The shell, natural size. . . . .   <i>c.</i> The sculpture, enlarged.	
Fig. 7. <i>Pleurotoma (Spirotropis) studeriana</i> , v. Mart. . . . .	322
(Copied from Professor v. Martens' Conchologische Mittheilungen, vol. i. pt. 3, pl. viii. fig. 3.)	
<i>a, b.</i> The shell, natural size. . . . .   <i>c.</i> The apex, enlarged.	
Fig. 8. <i>Pleurotoma (Thesbia) corpulenta</i> , n. sp. . . . .	331
<i>a.</i> The natural size. . . . .   <i>b.</i> The shell, enlarged.	
Fig. 9. <i>Pleurotoma (Spirotropis) bulbacea</i> , n. sp. . . . .	335
<i>a.</i> The natural size. . . . .   <i>c.</i> The operculum, magnified 10 diameters.	
<i>b.</i> The shell, enlarged. . . . .	
Fig. 10. <i>Pleurotoma (Surcula) hemimeres</i> , n. sp. . . . .	294
<i>a.</i> The natural size. . . . .   <i>b.</i> The shell, enlarged	
Fig. 11. <i>Pleurotoma (Thesbia) translucida</i> , n. sp. . . . .	330
<i>a.</i> The natural size. . . . .   <i>b.</i> The shell, enlarged.	