

8. *Rossia mölleri*, Steenstrup.1856. *Rossia Mölleri*, Stp., Hectocotyl., p. 198, pl. ii. fig. 1.9. *Rossia tenera* (Verrill) (p. 118).1880. *Heteroteuthis tenera*, VII., Amer. Journ. Sci. and Arts., vol. xx. p. 392.

1881. " VII., Ceph. N. E. Amer., p. 357, pl. xlvi. figs. 2, 3; pl. xlvii. fig. 5.

1883. " VII., " Blake Suppl., p. 111.

10. *Rossia patagonica*, E. A. Smith (Pl. XV. figs. 10–18; p. 119).1881. *Rossia patagonica*, E. A. Sm., Proc. Zool. Soc. Lond., p. 22, pl. iii. fig. 3.11. *Rossia megaptera*, Verrill.1881. *Rossia megaptera*, VII., Ceph. N. E. Amer., p. 349, pl. xxxviii. fig. 1; pl. xlvi. fig. 6.*Heteroteuthis*, Gray, 1849 (*non* Verrill).*Rossia*, Vérany et Auctt. (*pars*).1. *Heteroteuthis dispar* (Rüppell), Gray.1845. *Sepiola dispar*, Rüpp., Giorn. Gab. Messina, t. xxvi (*fide* Vér.)1849. *Rossia (Heteroteuthis) dispar*, Gray, B.M.C., p. 90.1851. " *dispar*, Vér., Céph. médit., p. 63, pl. xxiii. figs. d–h.

1857. " " Trosch., Archiv f. Naturgesch., Jahrg. xxiii. Bd. i. p. 62, pl. iv. figs. 7, 8.

1858. " " Claus, *Op. cit.*, Jahrg. xxiv. Bd. i. p. 259, Taf. x. fig. 5.*Promachoteuthis*, Hoyle, 1885.1. *Promachoteuthis megaptera*, Hoyle (Pl. XIV. figs. 10–14; p. 120).1885. *Promachoteuthis megaptera*, Hoyle, Narr. Chall. Exp., vol. i. p. 273, fig. 109.

1885. " " Hoyle, Diagnoses II., p. 182.

Family VIII. SEPIARII, Steenstrup, 1861.

SEPIDÆ, d'Orb. (*pars*).

Subfamily SEPIADARII, Steenstrup, 1881.

Sepioloidea, d'Orbigny, 1839.1. *Sepioloidea lineolata* (Quoy et Gaimard), d'Orbigny.1832. *Sepiola lineolata*, Q. et G., Voy. "Astrolabe," t. ii. p. 82, pl. v. figs. 8–13.1839. *Sepioloidea lineata*, d'Orb., Céph. acét., p. 240; Sépioles, pl. iii. figs. 10–18.

1881. " " Stp., Sepiadarium og Idiosepius, p. 233.

1884. " " Brock, Zeitschr. f. Wiss. Zool., Bd. xl. pp. 105–120.