

Indo-Pacific Species.

Pilumnus vespertilio (Fabricius) = *Pilumnus ursulus*, Adams and White; and
Pilumnus mus, Dana. Indo-Pacific Region (to 17 fathoms).

Pilumnus tomentosus, Latreille. Off Port Philip, Australia, 33 fathoms (Station 161); Bass Strait, 38 fathoms (Station 162).

Pilumnus peronii, Milne Edwards. Seas of Asia.

Pilumnus forskålii, Milne Edwards = *Cancer incanus*, Forskål. Coasts of Egypt.

This is possibly a variety of *Pilumnus vespertilio*, since Dr. Kossmann unites *Pilumnus ursulus* with this species, but in specimens from the Red Sea, referred to *Pilumnus forskålii* in the collection of the British Museum, not only are the tubercles of the palms of the chelipeds more distinctly longitudinally striate, but also the interior margin of the front is armed with longer spinules.

Pilumnus lanatus, Latreille. East Indies; North and East Australia, to 11 fathoms (H.M.S. "Alert"); Tasmania.

Pilumnus asper (Rüppell). Red Sea.

Pilumnus minutus, de Haan = *Pilumnus hirsutus*, Stimpson. Japan and Corean Straits (to 50 fathoms); North China Sea; New Caledonia; North-East Australia; Holborn Island (20 fathoms).

Pilumnus scabriusculus, Adams and White. Eastern Seas; Samboangan, Philippines.¹

Pilumnus globosus, Dana. Tahiti and Paumotu Islands; Japan (50 fathoms).

Pilumnus lævimanus, Dana. Balabac Passage; North of Borneo; New Caledonia.

Pilumnus lævis, Dana. Mangsi Islands, Straits of Balabac.

Pilumnus tenellus, Dana. Sooloo Sea, or Straits of Balabac.

Pilumnus rufopunctatus, Stimpson. East and South Australia.

Pilumnus fissifrons, Stimpson. East Australia.

Pilumnus forficigerus, Stimpson. Oosima, Japan (30 fathoms).

Pilumnus lapillimanus, Stimpson. North China Sea (25 fathoms).

Pilumnus marginatus, Stimpson. Loo-Choo.

Pilumnus dorsipes, Stimpson. Hong Kong (10 fathoms).

Pilumnus (?) pugilator (A. Milne Edwards). Lifu Island; North-East to East Australia (to 17 fathoms).

Pilumnus fragifer, A. Milne Edwards. Indian Ocean (perhaps a *Pilumnopeus*).

Pilumnus elatus, A. Milne Edwards. Upolu, Samoan Islands.

Pilumnus ovalis, A. Milne Edwards. Sandwich Islands.

¹ *Pilumnus dilatipes*, Adams and White, cannot be referred to this genus. Not only are the palatal ridges nearly obsolete, but the ambulatory legs are strongly cristated. It should probably be constituted the type of a distinct genus, *Lophopilumnus*.