

PLATE XII.

	Diam.	Page
Fig. 1. <i>Trapezia guttata</i> , Rüppell, var., adult female, . . . ×	3½	166
Fig. 1a. Chelipede, . . . . . ×	about 3	
Fig. 2. <i>Trapezia rufopunctata</i> (Herbst), var. <i>intermedia</i> , nov., adult female, . . . . . ×	2	168
Fig. 2a. Inferior view of the cervical region, showing the form of the antennæ, part of the orbits, and exterior maxillipedes, . . . ×	2½	
Fig. 2b. Chelipede, . . . . . ×	1½	
Fig. 3. <i>Pseudozius bouvieri</i> , var. <i>mellissii</i> , Miers, adult female, . . .	nat. size.	143
Fig. 3a. Inferior view of the cervical region, showing the form of the antennæ and exterior maxillipedes, . . . ×	about 2	
Fig. 3b. Exterior view of a chelipede, . . . . .	nat. size.	
Fig. 3c. Post-abdomen, . . . . . ×	1½	
Fig. 4. <i>Sphærozius nitidus</i> , Stimpson, adult male, . . . . . ×	about 5	144
Fig. 4a. Inferior view of the cervical region, showing the form of the antennæ and exterior maxillipedes, . . . ×	6	
Fig. 4b. Exterior view of a chelipede, . . . . . ×	4½	
Fig. 4c. Post-abdomen, . . . . . ×	8	