

Bermuda, Hungry Bay (swamps).

Eurytium limosum (Say).

Goniopsis cruentatus (Latreille).

Bahia, shallow water (7 to 20 fathoms).

Leptopodia sagittaria (Fabricius).

Metoporphis forficulatus, A. Milne
Edwards.

Notolopas brasiliensis, n. sp.

Picroceroides tubularis, n. gen.
and sp.

Macrocaloma trispinosa (Latreille).

Macrocaloma septemspinosa (Stimp-
son).

Macrocaloma concava, n. sp.

Mithrax cornutus, de Saussure.

Mithrax forceps, A. Milne Edwards.

Lambrus guérinii, F. de B. Capello.

Lambrus serratus, Milne Edwards.

Heterocrypta granulata (Gibbes).

Actæa rufopunctata, Milne Edwards,
var. *nodosa*, Stimpson.

Micropanope spinipes, A. Milne
Edwards.

Pilumnus brasiliensis, n. sp.

Pilumnus floridanus, Stimpson.

Pilumnus fragosus, A. Milne
Edwards.

Cronius bispinosus, n. sp.

Ocypoda arenaria (Catesby).

Sesarma mülleri, A. Milne Edwards.

Persephona punctata (Browne).

Lithadia cariosa, Stimpson, var.

Iliacantha intermedia, n. sp.

STATIONS 122-122c.—Off Barra Grande, 30 to 400 fathoms; lat. 9° 5' 0" to 9° 10' 0" S.,
long. 34° 49' 0" to 34° 53' 0" W.; bottom, red mud. Trawled.

Leptopodia sagittaria (Fabricius).

Metoporphis forficulatus, A. Milne
Edwards.

Podochela riisei (Stimpson).

Herbstia (*Herbstiella*) *depressa*,
Stimpson.

Mithrax hispidus (Herbst), var.
pleuracanthus, Stimpson.

Pilumnus floridanus, Stimpson.

Neptunus (*Hellenus*) *spinicarpus*,
Stimpson.

Bathyplox typhlus, var. *oculiferus*,
nov.

Fernando Noronha Island, shallow water (7 to 20 fathoms).

Apocremnus septemspinus, A.
Milne Edwards.

Picroceroides tubularis, n. gen.
and sp.

Macrocaloma concava, n. sp.

Mithrax coronatus (Herbst).

Mithrax sp.

Mithrax (*Mithraculus*) *sculptus*
(Lamarck).

Mithrax forceps, A. Milne Edwards.

Grapsus maculatus (Catesby).

Calappa gallus (Herbst).