

STATION 7. In the Faerøe Channel. August 12, 1880. Lat. $59^{\circ} 37' N.$, long. $7^{\circ} 19' W.$
Depth 530 fathoms.

Psilaster andromeda.

STATION 8. In the Faerøe Channel. August 17, 1880. Lat. $60^{\circ} 3' N.$, long. $5^{\circ} 51' W.$
Depth 540 fathoms. Ooze. Bottom temperature $29^{\circ} \cdot 2$ Fahr.; surface temperature $56^{\circ} \cdot 5$ Fahr.

Pontaster tenuispinus, var. *platynota.*

Plutonaster bifrons.

IV. "TRITON" EXPEDITION.

STATION 1. In the Faerøe Channel. August 4, 1882. Lat. $59^{\circ} 51' 30'' N.$, long. $6^{\circ} 21' 0'' W.$ Depth 240 fathoms. Bottom temperature $47^{\circ} \cdot 6$ Fahr.

Cribrella oculata. Also at Stations 10 and 11.

STATION 2. In the Faerøe Channel. August 5, 1882. Lat. $59^{\circ} 37' 30'' N.$, long. $6^{\circ} 49' 0'' W.$ Depth 530 fathoms. Bottom temperature $46^{\circ} \cdot 2$ Fahr.

Pteraster militaris.

STATION 3. In the Faerøe Channel. August 8, 1882. Lat. $60^{\circ} 39' 30'' N.$, long. $9^{\circ} 6' 0'' W.$ Depth 87 fathoms. Bottom temperature $49^{\circ} \cdot 5$ Fahr.

Luidia ciliaris.

Hippasteria plana.

STATION 5. In the Faerøe Channel. August 10, 1882. Lat. $60^{\circ} 11' 0''$ to $60^{\circ} 20' 0'' N.$, long. $8^{\circ} 15' 0''$ to $8^{\circ} 8' 0'' W.$ Depth 433 to 285 fathoms. Bottom temperature $43^{\circ} \cdot 5$ to $40^{\circ} \cdot 8$ Fahr.

Rhegaster murrayi.

Asterias (Leptasterias) mülleri.

STATION 9. In the Faerøe Channel. August 23, 1882. Lat. $60^{\circ} 5' 0'' N.$, long. $6^{\circ} 21' 0'' W.$ Depth 608 fathoms. Bottom temperature $30^{\circ} \cdot 0$ Fahr.

Pontaster tenuispinus, var. *platynota.*

Pteraster militaris, var. *prolata.*