

	Depth in Fathoms.	Occurrence in other Zones.
<i>ECHINASTERIDÆ.</i>		
<i>ECHINASTERINÆ.</i>		
<i>Echinaster spinulifer</i>	127
<i>HELIASTERIDÆ.</i>		
<i>Heliaster helianthus</i>
<i>PEDICELLASTERIDÆ.</i>		
<i>Pedicellaster scaber</i>	25
<i>Pedicellaster hypernotius</i>	140
<i>ASTERIIDÆ.</i>		
<i>Asterias meridionalis</i>	127
<i>Asterias perrieri</i>	110
<i>Asterias torquata</i>	25
<i>Asterias glomerata</i>	55
<i>Asterias versicolor</i>	50
<i>Asterias amurensis</i>	25
<i>Asterias cunninghami</i>	55
<i>Asterias (Cosinasterias) tomidata</i>	45
<i>Asterias (Smilasterias) scalprifera</i>	75
<i>Asterias (Smilasterias) trivenis</i>	150
<i>Asterias (Leptasterias) compta</i>	85
<i>Asterias (Stolasterias) gemmifera</i>
<i>Asterias (Stolasterias) tenuispina</i>
<i>Asterias (Stolasterias) calamaria</i>	40
<i>Asterias (Stolasterias) volsellata</i>	95
<i>Asterias (Stolasterias) eustyla</i>	150
<i>Asterias (Stolasterias) africana</i>	20
<i>Calvasterias stolidota</i>	10
<i>BRISINGIDÆ.</i>		
<i>Labidiaster radiusus</i>	55
<i>Labidiaster annulatus</i>	150	Also Abyssal.

The above table comprises 141 species and varieties found in the Littoral Zone. Of these 8 extend into the Continental Zone, and 9 extend into the Abyssal Zone.

(2.) CONTINENTAL ZONE.

Species found in depths between 150 fathoms and 500 fathoms.

	Depth in Fathoms.	Occurrence in other Zones.
<i>ARCHASTERIDÆ.</i>		
<i>PARARCHASTERINÆ.</i>		
<i>Pararchaster spinosissimus</i>	425
<i>Pontaster planeta</i>	245
<i>Pontaster oxyacanthus</i>	345
<i>PLUTONASTERINÆ.</i>		
<i>Plutonaster marginatus</i>	425
<i>ARCHASTERINÆ.</i>		
<i>Archaster typicus</i>	250	Also Littoral.