

	Depth in Fathoms.	Occurrence in other Zones.
<i>SOLASTERIDÆ.</i>		
<i>SOLASTERINÆ.</i>		
<i>Crossaster papposus</i>	384	Also Abyssal.
<i>Lophaster furcifer</i>	384	Also Abyssal.
<i>ECHINASTERIDÆ.</i>		
<i>ECHINASTERINÆ.</i>		
<i>Cribrella oculata</i>	458	Also Littoral.
<i>ASTERIIDÆ.</i>		
<i>Asterias (Leptasterias) mülleri</i>	312	Also Littoral.
<i>BRISINGIDÆ.</i>		
<i>Odinia pandina</i>	440
<i>Brisinga endecacnemos</i>	469	Also Abyssal.
<i>Brisinga coronata</i>	458	Also Abyssal.

(3.) ABYSSAL ZONE.

Species found in depths greater than 500 fathoms.

	At or greater than						Greatest Depth recorded.	Occurrence in other Zones.
	500	750	1000	1500	2000	2500		
<i>ARCHASTERIDÆ.</i>								
<i>PLUTONASTERINÆ.</i>								
<i>Plutonaster bifrons</i>	+	...	+	1360
<i>Plutonaster (Tethyaster) parelii</i>	+	1360
<i>PORCELLANASTERIDÆ.</i>								
<i>CTENODISCINÆ.</i>								
<i>Ctenodiscus corniculatus</i>	+	632	Also Continental.
<i>ASTROPECTINIDÆ.</i>								
<i>ASTROPECTININÆ.</i>								
<i>Leptoptychaster arcticus</i>	+	690	Also Continental.
<i>Psilaster andromeda</i>	+	690	Also Continental.
<i>GYMNASTERIIDÆ.</i>								
<i>Marginaster fimbriatus</i>	+	1360
<i>Lasiaster villosus</i>	+	542
<i>ZOROASTERIDÆ.</i>								
<i>Zoroaster fulgens</i>	+	767
<i>SOLASTERIDÆ.</i>								
<i>SOLASTERINÆ.</i>								
<i>Crossaster papposus</i>	+	640	Also Continental.
<i>Lophaster furcifer</i>	+	605	Also Continental.
<i>KORETHRASTERINÆ.</i>								
<i>Korethraster hispidus</i>	+	632
<i>PTERASTERIDÆ.</i>								
<i>PTERASTERINÆ.</i>								
<i>Hymenaster pellucidus</i>	+	580
<i>BRISINGIDÆ.</i>								
<i>Brisinga endecacnemos</i>	+	...	+	1095	Also Continental.
<i>Brisinga coronata</i>	+	630	Also Continental.

The following species were also collected by the "Porcupine," but the depth is not recorded:—

*Palinipes membranaceus.**Chelaster longipes.**Stichaster roseus.*