

	LOCALITIES.	RANGE IN DEPTH : FATHOMS.	NATURE OF THE SEA-BOTTOM.	
PHANEROZONIA.				
<i>ARCHASTERIDÆ.</i>				
<i>ARCHASTERINÆ.</i>				
<i>Archaster</i> , Müller and Troschel, <i>emend.</i>				
* <i>A. typicus</i> , Müller and Troschel	Eastern Archipelago; Pelew Islands; New Guinea; Port Darwin; Port Essington; Cape Grenville; Port Denison; New Caledonia; Fiji and Tonga Islands; Nicobar Islands; Andaman Islands; Mergui; Japan (!).	Shallow to 250	Coral reefs; Green mud.	(1)
<i>Archaster non sensu stricto</i> :—				
<i>A. americanus</i> , Verrill	E. coast of N. America.	45-225	(2)
<i>A. bairdii</i> , Verrill	E. coast of N. America.	351-396	(3)
<i>A. efflorescens</i> , Perrier	N. of Havana.	955	(4)
<i>A. floreæ</i> , Verrill ¹	E. coast of N. America.	86-410	(5)
<i>A. formosus</i> , Verrill	E. coast of N. America.	1467-1608	(6)
<i>A. insignis</i> , Perrier	N. of Havana, W. of Tortugas.	1930	(7)
<i>A. robustus</i> , Verrill	E. coast of N. America.	938-1467	(8)
<i>A. sepitus</i> , Verrill	E. coast of N. America.	368-858	(9)
<i>ARCHASTERIDÆ incertæ sedis</i> :				
<i>Benthopecten</i> , Verrill.				
<i>B. spinosus</i> , Verrill	E. coast of N. America.	855-1917	(10)
<i>Blakiaster</i> , Perrier.				
<i>B. conicus</i> , Perrier	Off Grenada; off Havana.	92-175	(11)
<i>Cheiraster</i> , Studer.				
<i>C. gazellæ</i> , Studer	N.W. Australia.	195	Grey mud.	(12)
<i>C. pedicellaris</i> , Studer	Off E. coast of Australia.	550	Globigerina ooze.	(13)
<i>C. (?) folini</i> , Perrier	"Talisman" dredgings.	(14)

¹ Probably either *Bathyastrer* or *Psilaster* (!).