

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
PHANEROZONIA				
<i>PENTAGONASTERIDÆ.</i>				
Subfamily GONIODISCINÆ, Sladen.				
<i>Stellaster</i> , Gray.				
<i>S. equestris</i> (Retzius), Müller and Troschel	China; Japan; Formosa; Port Darwin; Cape Cleveland.	(1)
<i>S. granulatus</i> , Perrier, sp.	New South Wales.	(2)
* <i>S. incei</i> , Gray	{ Philippine Islands; Amboina; Sumatra; (?) Corean Strait; New Guinea; Arafura Sea; Torres Strait; N., N.E., and S. Australia.	4-50	{ Coral mud; Green mud; Blue mud. }	(3)
* <i>S. princeps</i> , Sladen	Torres Strait.	6	Coral mud.	(4)
<i>S. squamulosus</i> , Studer	N.W. Australia.	60	(5)
<i>S. tuberculatus</i> , von Martens	China (<i>vide</i> Perrier).	(6)
<i>Ogmaster</i> , von Martens.				
<i>O. capella</i> (Müller and Troschel), von Martens	(?) China or Japan.	(7)
<i>Leptogonaster</i> , Sladen.				
* <i>L. cristatus</i> , Sladen	Off the Philippine Islands.	100-115	Green mud.	(8)
<i>Goniodiscus</i> , Müller and Troschel, emend. Perrier.				
<i>G. articulatus</i> (Linné), de Loriol	{ Singapore; Mergui Archipelago.	{ Sub-littoral.	(9)
<i>G. cuspidatus</i> (Lamarck), Müller and Troschel	{ "Mers australes" (Peron and Lesueur).	{ ?	(10)
<i>G. forficulatus</i> , Perrier	Migupou.	(11)
<i>G. gracilis</i> , Gray	Moluccas.	(12)
<i>G. granulosus</i> (Gray), Perrier	China (<i>vide</i> Perrier).	(13)