

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
PHANEROZONIA.				
<i>ANTHENEIDÆ.</i>				
<i>Anthenea</i> , Gray.				
<i>A. articulata</i> , Perrier	{ Seychelle Islands; China (<i>fide</i> Perrier). }	(1)
<i>A. flavescens</i> (Gray), Perrier {	Port Jackson; Freemantle, W. Australia (<i>fide</i> Bell). }	0-5	(2)
<i>A. grayi</i> , Perrier	?	?	?	(3)
<i>A. pentagonula</i> (Lamarck), Perrier {	Hong Kong; Madras (<i>fide</i> Bell); N.W. Australia (<i>fide</i> Studer). }	5	(4)
* <i>A. tuberculosa</i> , Gray	{ Port Essington (N. Australia), Torres Strait; Port Jackson (<i>fide</i> Bell). }	6 ...	Coral mud.	(5)
<i>A. (?) spinulosa</i> , Gray, sp.	{ Philippine Islands; Timor; Fiji Islands; N.E. Australia. }	...	Coral reefs.	(6)
<i>Goniaster</i> (Agassiz), <i>emend.</i> Perrier.				
<i>G. obtusangulus</i> (Lamarck), Perrier	Indian Ocean (?).	(7)
<i>Hippasteria</i> , Gray.				
* <i>H. plana</i> (Linck), Gray	{ Off the British, Scandinavian, and Murman coasts; off E. of North America, N. of Cape Cod. }	30-150	Clay, gravel, stones.	(8)
Family <i>PENTACEROTIDÆ</i> , Gray, <i>emend.</i> Perrier.				
<i>Pentaceros</i> , Linck.				
<i>P. affinis</i> , Müller and Troschel, sp.	India.	(9)
<i>P. alveolatus</i> , Perrier	New Caledonia.	(10)
<i>P. australis</i> , Lütken, sp.	Australia.	(11)