

	LOCALITIES.	RANGE IN DEPTH. FATHOMS.	NATURE OF THE SEA- BOTTOM.	
PHANEROZONIA. <i>PENTACEROTIDÆ.</i>				
<i>Pentaceros</i> , Linck.				
<i>P. valvulatus</i> , Müller and Troschel	S.W. Australia.	(1)
<i>P. verrucosus</i> , Müller and Troschel, sp.	India.	(2)
<i>P. westermannii</i> , Lütken, sp.	Bengal.	(3)
<i>Nidorellia</i> , Gray.				
<i>N. armata</i> (Gray), Verrill.	W. coast of Central America, extending from California to Ecuador.	(4)
<i>N. michelini</i> , Perrier.	Mazatlan.	(5)
<i>Amphiaster</i> , Verrill.				
<i>A. insignis</i> , Verrill.	La Paz.	(6)
<i>Pentaceropsis</i> , Sladen.				
* <i>P. obtusatus</i> , Bory de Saint Vincent, sp.	Philippine Islands; Mauri- tius (?).	Shallow.	(7)
<i>Culcita</i> , Agassiz.				
<i>C. acutispina</i> , Bell.	New Hebrides.	(8)
<i>C. arenosa</i> (Val.), Perrier.	Sandwich Islands.	(9)
<i>C. coriacea</i> , Müller and Troschel.	Red Sea; Mozambique; Mau- ritius.	(10)
<i>C. grec</i> , Müller and Troschel.	Molucca; Andaman Islands (<i>sic</i> Bell).	(11)
* <i>C. novae-guineæ</i> , Müller and Troschel.	New Guinea; New Hanover; N.E. Australia; Nicobar Islands (<i>sic</i> Lütken).	10	Sand.	(12)
<i>C. pentangularis</i> , Gray.	Mozambique; Torres Strait; Fiji Islands; N.E. and W. Australia.	(13)