

| | LOCALITIES. | RANGE IN DEPTH: FATHOMS. | NATURE OF THE SEA-BOTTOM. | |
|--|--|--------------------------|---|------|
| PHANEROZONIA. | | | | |
| <i>Gymnasteriidae.</i> | | | | |
| <i>Gymnasteria</i> , Gray. | | | | |
| (1) <i>G. biserrata</i> , von Martens | Larentuka. | ... | | (1) |
| <i>G. valvulata</i> , Perrier | Galapagos Islands. | ... | | (2) |
| <i>Tylaster</i> , Danielssen and Koren. | | | | |
| <i>T. willei</i> , Danielssen and Koren { | Off Spitzbergen, and between Spitzbergen and Norway. | } 416-1200 { | Clay; Biloculina clay; Bluish green clay. } | (3) |
| <i>Porania</i> , Gray. | | | | |
| * <i>P. antarctica</i> , Smith | Kerguelen; Marion Island; Prince Edward Island; Crozet Islands; South Georgia. | } 50-1600 { | Diatom ooze (1600 fathoms). | (4) |
| * <i>P. glaber</i> , Sladen | Kerguelen Island. | 30-127 | | (5) |
| <i>P. grandis</i> , Verrill | E. of North America. | 66-373 | | (6) |
| * <i>P. magellanica</i> , Studer | Off Patagonia; in Strait of Magellan. | } 45 | Green sand. | (7) |
| <i>P. pulvillus</i> (O. F. Müller), Norman | Off the coasts of Scandinavia and Britain. | } 15-106 | | (8) |
| * <i>P. spiculata</i> , Sladen | Off Heard Island; off the Arrou Islands. | } 75-800 { | Coarse gravel; Volcanic mud; Green mud. | (9) |
| <i>Marginaster</i> , Perrier. | | | | |
| <i>M. echinulatus</i> , Perrier | Off Barbados. | 69 | | (10) |
| <i>M. fimbriatus</i> , Sladen | N. of Ireland. | 1360 | | (11) |
| <i>M. pectinatus</i> , Perrier | Off coast of Yucatan. | 95 | | (12) |
| <i>M. pentagonus</i> , Perrier | "Talisman" dredging, No. 37. | 218 | | (13) |
| <i>M.</i> (1) <i>caprensensis</i> , Gasco, sp. | Off Island of Capri. | 54-82 | | (14) |