

	LOCALITIES.	RANGE IN DEPTH : FATHOMS.	NATURE OF THE SEA-BOTTOM.	
PHANEROZONIA.				
<i>ASTERINIDÆ.</i>				
<i>ASTERININÆ.</i>				
<i>Asterina</i> , Nardo.	Cape of Good Hope; Natal; Madagascar; Mauritius; Java; Moluccas; Philippine Islands; New Guinea; S. and E. Australia; St. Paul Island.	10	Coral reefs, &c.	(1)
* <i>A. exigua</i> (Lamarck), Perrier				
<i>A. fimbriata</i> , Perrier . . .	Island of Bourbon; Strait of Magellan.	(2)
* <i>A. folium</i> (Lütken), Agassiz .	Antilles; Bermuda.	(3)
<i>A. gayi</i> , Perrier	Chili.	(4)
<i>A. gibbosa</i> (Pennant), Forbes	Off European and N. African coasts; Mediterranean; Azores (<i>vide</i> Barrois). Also reputed to occur in the Indian Ocean and Eastern Archipelago; but probably <i>A. cepheus</i> is the species in question.	0-35	(5)
<i>A. granifera</i> (Gray), Perrier .	Cape of Good Hope.	(6)
<i>A. granulosa</i> , Perrier. . . .	Sandwich Islands.	(7)
* <i>A. gunnii</i> , Gray	Cape of Good Hope; Port Mollo; South Australia; Port Phillip; Tasmania.	7	(8)
<i>A. lymani</i> , Perrier	Barbados.	120-140	Broken shells and corals.	(9)
<i>A. marginata</i> (Val.), Perrier .	Rio Janeiro; Senegal; (?) Canary Islands.	(10)
<i>A. miniata</i> (Brandt), Perrier .	Sitcha; California.	(11)
<i>A. minuta</i> , Gray	Antilles.	(12)
<i>A. modesta</i> (Verrill), Perrier .	Panama; Pearl Islands.	(13)
<i>A. novæ-zealandiæ</i> , Perrier . .	New Zealand.	(14)