

| | LOCALITIES | RANGE IN DEPTH: FATHOMS. | NATURE OF THE SEA-BOTTOM. | |
|--|--|--------------------------|--------------------------------|------|
| PHANEROZONIA. | | | | |
| <i>ASTERINIDÆ.</i> | | | | |
| <i>ASTERININÆ.</i> | | | | |
| <i>Asterina</i> , Nardo. | | | | |
| <i>A. obtusa</i> (Gray), Perrier . . . | Panama. | 1-2 | | (1) |
| * <i>A. pectinifera</i> (Müller and Troschel), von Martens . . . | Japan. | 5-25 | | (2) |
| * <i>A. penicillaris</i> (Lamarck), von Martens | Red Sea; Port Natal; Java; Flores; Moluccas; Australia; Japan. | 8-50 | | (3) |
| <i>A. pilosa</i> , Perrier | Dominica. | 118 | } Fine sand and broken shells. | (4) |
| <i>A. pusilla</i> , Perrier | Talcahuano (Chili). | ... | | (5) |
| <i>A. pygmaea</i> , Verrill | E. of North America. | ... | | (6) |
| * <i>A. regularis</i> , Verrill | Port Mollo; Port Jackson; New Zealand. | 10 | | (7) |
| <i>A. setacea</i> (Val.), Perrier | ? | ? | ? | (8) |
| <i>A. squamata</i> (Val.), Perrier | Senegal. | ... | | (9) |
| <i>A. stellaris</i> , Perrier | ? | ? | ? | (10) |
| <i>A. trochiscus</i> (Retzius), Perrier | Indian Ocean (?) | ... | | (11) |
| <i>A. uega</i> (Val.), Perrier | Red Sea; Mauritius. | ... | | (12) |
| <i>Disasterina</i> , Perrier. | | | | |
| <i>D. abnormalis</i> , Perrier | New Caledonia. | ... | | (13) |
| <i>D. ceylanica</i> , Döderlein | Ceylon. | ... | | (14) |
| Subfamily PALMIPEDINÆ, Sladen. | | | | |
| <i>Palmipes</i> , Linck. | | | | |
| * <i>P. diaphanus</i> , Sladen | N. of Admiralty Island. | 150 | Coral mud. | (15) |