

	LOCALITIES.	RANGE IN DEPTH: FATHOMS.	NATURE OF THE SEA-BOTTOM.	
<b>PHANEROZONIA</b>				
<i>ASTERINIDÆ.</i>				
<i>PALMIPEDINÆ.</i>				
<i>Palmipes</i> , Linck.				
<i>P. membranaceus</i> , Linck . . .	{ British Isles; English Channel; coast of France; Mediterranean; Adriatic.	} 20-100	.....	(1)
<i>P. rosaceus</i> (Lamarck), Dujardin and Hupe . . . . .	{ Japan; Bay of Bengal.	} ...	.....	(2)
<i>Stegnaster</i> , Sladen.				
<i>S. inflatus</i> , Hutton, sp. . . .	New Zealand.	...	.....	(3)
<i>S. wesseli</i> , Perrier, sp. . . .	Barbados; off Florida.	101	.....	(4)
<i>ASTERINIDÆ incertæ sedis:</i>				
<i>Tremaster</i> , Verrill.				
<i>T. mirabilis</i> , Verrill . . . . .	{ Off George's Bank, Nova Scotia, and Newfoundland.	} 150-250	.....	(5)
Order CRYPTOZONIA, Sladen.				
Family LINCKIIDÆ, Perrier, emend.				
Subfamily CHETASTERINÆ, Sladen.				
<i>Chetaster</i> , Müller and Troschel.				
* <i>C. longipes</i> (Retzius), Sars . . .	{ Mediterranean; Azores; Bermuda; S. of Cape Palmas ( <i>vide</i> Studer).	} 30-450	Coral; Volcanic mud.	(6)
<i>C. nodosus</i> , Perrier . . . . .	{ Guadeloupe; Cape Verde Islands ( <i>vide</i> Studer). }	} ...	.....	(7)
(!) <i>C. hermanni</i> , Müller and Troschel . . . . .	{ ?	} ?	?	(8)