

	LOCALITIES.	RANGE IN DEPTH : FATHOMS.	NATURE OF THE SEA-BOTTOM.	
<b>CRYPTOZONIA.</b>				
<i>LINCKIIDÆ.</i>				
Subfamily LINCKIINÆ, Sladen.				
<i>Fromia</i> , Gray.				
<i>F. balansæ</i> , Perrier . . . .	New Caledonia.	...	.....	(1)
<i>F. indica</i> , Perrier . . . .	{ Indian Ocean; Andaman Islands.	}	.....	(2)
<i>F. japonica</i> , Perrier . . . .				
* <i>F. milleporella</i> (Lamarck), Gray	{ Red Sea; Mauritius; Madagascar; Ceylon; Moluccas; Amboina; New Caledonia; Fiji, Samoa, and Loo Choo Islands.	}	Shallow.	Coral reefs.
<i>F. monilis</i> (Val.), Perrier . . . .				
<i>F. tumida</i> , Bell . . . .	Ceylon; Andaman Islands.	...	.....	(6)
<i>Ferdina</i> , Gray.				
<i>F. cancellata</i> , Grube, sp. . . .	{ Fiji Islands (Muss. Copenhagen and Berlin). }	}	.....	(7)
<i>F. cumingii</i> , Gray . . . .				
<i>F. flavescens</i> , Gray . . . .	Mauritius.	...	.....	(9)
<i>F. küllii</i> , Müller and Troschel, sp.	Java.	...	.....	(10)
<i>Oplidiaster</i> , Agassiz.				
<i>O. agassizii</i> , Perrier . . . .	Juan Fernandez.	...	.....	(11)
<i>O. arenatus</i> (Lamarck), Perrier . . . .	?	?	?	(12)
.....	...	...	.....	)
* <i>O. attenuatus</i> , Gray . . . .	Mediterranean; Azores.	450	Volcanic mud.	(13)
<i>O. bicolor</i> (Lamarck), Perrier . . . .	?	?	?	(14)