

	GEOGRAPHICAL AREAS.							BATHYMETRICAL ZONES.			BOTTOM TEMPERATURE FAHRENHEIT.	SYNONYMA AND REMARKS.
	North Atlantic.	South Atlantic.	Indian Ocean.	Southern Ocean.	Eastern Archipelago.	North Pacific.	South Pacific.	Littoral, 0-100 Fathoms.	Oestental, 100-500 Fathoms.	Absyal, greater than 500 Fathoms.		
	I.	II.	III.	IV.	V.	VI.	VII.	L	C.	A.		
(1)	+	L	C	A	35°0-47°0	= <i>Pentadactylaster oculatus</i> , Liack; <i>Asterias oculata</i> , Pennant; <i>Linkia oculata</i> , Forbes (olim); <i>Henricia oculata</i> , Gray; <i>Echinaster oculatus</i> , Müller and Troschel. Incl.: <i>Asterias sanguinolenta</i> , O. F. Müller; <i>Asterias pertusa</i> , O. F. Müller; <i>Asterias spongiosa</i> , Fabricius; <i>Asterias seposita</i> , Retzius; <i>Echinaster Echrichii</i> , Müller and Troschel; <i>Echinaster sanguinolentus</i> , Sars; <i>Echinaster Sarnii</i> , Müller and Troschel; <i>Linkia pertusa</i> , Simpson; <i>Cribrella sanguinolenta</i> , Lütken; <i>Oribella Echrichii</i> , Dujardin and Hupé.
(2)	+	A	45°5-46°0	= <i>Cribrella oculata</i> , var. <i>cylindrella</i> , Bladen (sde Norman).
(3)	...	+	+	L	= <i>Echinaster (Oribella) ornatus</i> , Perrier (olim).
(4)	...	+	L
(5)	+	C
(6)	+	L	C	...	{ 46°76- 61°75 }
(7)	...	+	...	+	L	C
(8)	+	L
(9)	+	A	43°0
(10)	+	L
(11)	+	L
(12)	...	+	L	{ = <i>Othilia multispinosa</i> , Gray; <i>Cribella brasiliensis</i> , Dujardin and Hupé; <i>Othilia brasiliensis</i> , Agassiz.