

	LOCALITIES.	RANGE IN DEPTH : FATHOMS.	NATURE OF THE SEA-BOTTOM.	
CRYPTOZONIA.				
<i>BRISINGIDÆ.</i>				
<i>Freyella</i> , Perrier.				
* <i>F. polycnema</i> , Sladen . . .	N.E. of Kermadec Islands.	600	Hard ground.	(1)
* <i>F. remex</i> , Sladen . . .	S.E. of New Guinea.	2440	Red clay.	(2)
<i>F. sexradiata</i> , Perrier . . .	"Talisman" dredgings.	2733	(3)
<i>F. spinosa</i> , Perrier . . .	Off coast of the Soudan.	784	(4)
* <i>F. tuberculata</i> , Sladen . . .	{ Between the Canary Islands and Cape Verde Islands; between coast of Africa and Island of Ascension.	{ 2350-2400	Globigerina ooze.	(5)
<i>Colpaster</i> , Sladen.				
* <i>C. scutigerula</i> , Sladen . . .	S.W. of Canary Islands.	1525	Hard ground.	(6)
Genera of doubtful value :				
<i>Brisingaster</i> , de Loriol.				
<i>B. robillardi</i> , de Loriol . . .	Mauritius.	60	(7)
<i>Hymenodiscus</i> , Perrier.				
<i>H. agassizii</i> , Perrier . . .	Dominica ; Santa Cruz.	391-450	Sand and Grey mud.	(8)
<i>Gymnobrisinga</i> , Studer.				
<i>G. sarsii</i> , Studer . . .	{ N. or N.W. of Kerguelen Island.	{ 115 or 150	(9)

137 genera, 810 species, and 22 varieties are enumerated in this list.