

B. BATHYMETRICAL RANGE OF THE SPECIES.

I. Challenger Expedition—		PAGE
(1.) Littoral Zone		687
(3.) Continental Zone		690
(3.) Abyssal Zone		692
II. "Porcupine" Expeditions—		
(1.) Littoral Zone		695
(2.) Continental Zone		695
(3.) Abyssal Zone		696
III. "Knight Errant" Expedition—		
(1.) Littoral Zone		697
(2.) Continental Zone		697
(3.) Abyssal Zone		697
IV. "Triton" Expedition—		
(1.) Littoral Zone		698
(2.) Continental Zone		698
(3.) Abyssal Zone		698
Table showing the known Bathymetrical Range of the Genera of Asteroidea collected by the Challenger in each of the great Oceanic Areas		699

C. NATURE OF THE SEA-BOTTOM ON WHICH THE ASTEROIDEA COLLECTED BY THE CHALLENGER WERE FOUND, WITH AN ENUMERATION OF THE SPECIES OCCURRING ON EACH DEPOSIT.

I. Terrigenous Deposits—		
a. Gravel, Stones		703
b. Coarse Gravel		703
c. Sand		703
d. Sand, Gravel		704
e. Sand and Shells		704
f. Rock		704
g. Hard Ground, "Gravel, Shells"		705
h. Hard Ground		705
i. Mud		705
j. Blue Mud		706
k. Green Mud		707
l. Green Sand		709
m. Red Mud		709
n. Volcanic Mud		709
o. Greenish Volcanic Mud		710
p. Volcanic Sand		711
q. Coral Mud		711
r. Coral Reefs		712