

PLATE CI

Alpheus avarus (p. 544).

Fig. 1. Lateral view ; enlarged twice.

Alpheus neptunus (p. 563).

- „ 2. Lateral view ; enlarged twice.
- „ 2*k*. First pereopod ; larger chela, left.
- „ 2*k'*. First pereopod ; smaller chela, right.

Alpheus gracilipes (p. 561).

- „ 3. Lateral view ; enlarged three times.
- „ 3*c*. Cephalon, frontal region.
- „ 3*k*. First pereopod ; larger chela.
- „ 3*m*. Third pereopod ; terminal joints.
- „ 3*z*. Telson.

Alpheus biunguiculatus (p. 562).

- „ 4. Lateral view ; enlarged three times.
- „ 4*o*. Fifth pereopod ; terminal joints.
- „ 4*z*. Telson.

Betæus malleodigitus (p. 565).

- „ 5. Lateral view ; enlarged three times.
- „ 5*c*. Cephalon, frontal region.
- „ 5*c*. Second antennæ ; scaphocerite.
- „ 5*k*. First pereopod ; larger chela.
- „ 5*l*. Second pereopod.
- „ 5*p*. First pleopod, with parasites attached to peduncle ; stylamblys detached and enlarged.
- „ 5*par*. Parasitic vesicle, detached ; escaped ovum, enlarged.
- „ 5*z*. Telson.

Betæus microstylus (p. 566).

- „ 6. Lateral view ; enlarged three times.
- „ 6*c*. Cephalon, frontal region and scaphocerites.
- „ 6*c*. Second antennæ ; scaphocerite, enlarged.
- „ 6*z*. Telson.