

STATION 104. August 23, 1873; lat. $2^{\circ} 25'$ N., long. $20^{\circ} 1'$ W.; depth, 2500 fathoms; bottom, Globigerina ooze; bottom temperature, $36^{\circ}.6$. Trawled.

<i>Haliporus lœvis</i> (2 ♀).		<i>Acanthephyra kingsleyi</i> (1 ♂).
		<i>Hymenodora mollicutis</i> (1 ♀).

STATION 106. August 25, 1873; lat. $1^{\circ} 47'$ N., long. $24^{\circ} 26'$ W.; depth, 1850 fathoms; bottom, Globigerina ooze; bottom temperature, $36^{\circ}.6$. Trawled.

<i>Haliporus lœvis</i> (1 ♂).		<i>Sergestes oculatus</i> (4).
<i>Gennadas intermedius</i> (1).		<i>Sergestes longispinus</i> (1).
		<i>Acanthephyra acanthitelsonis</i> (1 ♂).

Tropical Atlantic. Surface.

<i>Lucifer typus.</i>		<i>Sergestes parvidens.</i>
<i>Lucifer reynaudii.</i>		<i>Peteinura gubernata</i> , at night.

STATION 107. August 26, 1873; lat. $1^{\circ} 22'$ N., long. $26^{\circ} 36'$ W.; depth, 1500 fathoms; bottom, Globigerina ooze; bottom temperature, $37^{\circ}.9$. Trawled.

<i>Acanthephyra acanthitelsonis</i> (1 ♂).		<i>Acanthephyra brevirostratis</i> (2).
--	--	---

St. Paul's Rocks. Surface.

<i>Panulirus guttatus</i> (3 ♀ 1 ♂).		<i>Lucifer reynaudii.</i>
<i>Lucifer typus.</i>		<i>Alpheus minus</i> (2 ♀).
		<i>Sergestes oculatus</i> (6).

STATION 113A. September 2, 1873; lat. $3^{\circ} 47' 0''$ S., long. $32^{\circ} 24' 30''$ W.; off Fernando Noronha; depth, 7 to 25 fathoms; bottom, volcanic sand and gravel.

<i>Artemesia longinaris</i> (1).		<i>Alpheus minus</i> (1 ♀).
----------------------------------	--	-----------------------------

STATION 120. September 9, 1873; lat. $8^{\circ} 37'$ S., long. $34^{\circ} 28'$ W.; off Pernambuco; depth, 675 fathoms; bottom, red mud. Trawled.

<i>Gennadus parvus</i> (1 ♀).		<i>Notostomus brevirostris</i> (1).
<i>Glyphocrangon aculeata</i> (1 ♀).		<i>Hymenodora mollis</i> (1 ♀).

STATION 122. September 10, 1873; lat. $9^{\circ} 5'$ S., long. $34^{\circ} 50'$ W.; off Barra Grande; depth, 350 fathoms; bottom, red mud. Trawled.

<i>Amphiplectus depressus</i> (2 ♀).		<i>Plesionika semilævis</i> (16).
<i>Plesionika uniproducta</i> (2 ♂ ♀).		<i>Nothocaris geniculatus</i> (17 ♀ ♂).
		<i>Campylonotus capensis</i> (2 ♂).