

STATION 126. September 12, 1873; lat.  $10^{\circ} 46'$  S., long.  $36^{\circ} 8'$  W.; depth, 770 fathoms; bottom, red mud. Trawled.

*Acanthephyra edwardsii* (2 ♂ ♀)

Off Bahia; from 7 to 22 fathoms.

<i>Sicyonia carinata.</i>		<i>Alpheus minus</i> (3 ♀ ♂).
		<i>Alpheus intrinsecus</i> (1 ♀).

South Atlantic, October 5, 1873; about lat.  $28^{\circ}$  S., long.  $27^{\circ}$  W. Surface.

*Sergestes longicollus.*

STATION 133. October 11, 1873; lat.  $35^{\circ} 41'$  S., long.  $20^{\circ} 55'$  W.; near Tristan da Cunha; depth, 1900 fathoms; bottom, Globigerina ooze; bottom temperature,  $35^{\circ}.4$ . Trawled.

<i>Willemoesia leptodactyla</i> (1 ♀).		<i>Benthesicymus altus</i> (3 ♂).
<i>Hemipenaeus spinidorsalis</i> (2).		<i>Benthesicymus mollis</i> (1).
<i>Aristeus armatus</i> (1 ♂).		<i>Pontophilus gracilis</i> (1 ♀).
<i>Benthesicymus iridescens</i> (2 ♀ ♂).		<i>Notostomus murrayi</i> (1).
		<i>Hymenodora mollicutis</i> (2 ♂ ♀).

STATION 135C. October 17, 1873; lat.  $37^{\circ} 25' 30''$  S., long.  $12^{\circ} 28' 30''$  W.; Nightingale Island, Tristan da Cunha; depth, 110 fathoms. Dredged.

*Palinosytus lalandii* (1).

STATION 137. October 23, 1873; lat.  $35^{\circ} 59'$  S., long.  $1^{\circ} 34'$  E.; depth, 2550 fathoms; bottom, red clay; bottom temperature,  $34^{\circ}.5$ . Dredged.

<i>Gennadas intermedius</i> (1 ♂).		<i>Sergestes profundus</i> (1 ♀).
------------------------------------	--	-----------------------------------

STATION 142. December 18, 1873; lat.  $35^{\circ} 4'$  S., long.  $18^{\circ} 37'$  E.; Agulhas Bank; depth, 150 fathoms; bottom, green sand; bottom temperature,  $47^{\circ}$ . Dredged.

<i>Merhippolyte agulhasensis</i> (5 ♂ ♀).		<i>Pandalus modestus</i> (3).
		<i>Chlorotocus incertus</i> (1).

STATION 144A. December 26, 1873; lat.  $46^{\circ} 48' 0''$  S., long.  $37^{\circ} 49' 30''$  E.; off Marion Island; depth, 69 fathoms; bottom, volcanic sand. Dredged.

*Nauticaris marionis* (67 ♀ ♂).