

Fiji Islands. Surface.

Penæus canaliculatus (many ♀ ♂).
Lucifer reynaudii.
Thalassocaris danæ (1). August
 11, 1874.

Thalassocaris stimpsoni (2). August
 11, 1874.
Oodeopus duplex.
Oodeopus serratus.

Off Kandavu, Fiji Islands.

Thalassina scorpionoides.
Stenopus hispidus.

Caricyphus serratarginis (1).
 August 11, 1874.

Kandavu. Fresh-water.

Bithynis lar (23 ♂ ♀).

Ovalau, Fiji Islands. Fresh-water rivers.

Bithynis lar.

STATION 175. August 12, 1874; lat. 19° 2' S., long. 177° 10' E.; Fiji Islands; depth, 1350 fathoms; bottom, Globigerina ooze; bottom temperature, 36°. Trawled.

Glyphocrangon acuminata (22 ♀ ♂).

STATION 176. August 15, 1874; lat. 18° 30' S., long. 173° 52' E.; off the New Hebrides.

Nematocarcinus productus.

STATION 177. August 18, 1874; lat. 16° 45' S., long. 168° 7' E.; off the New Hebrides; depth, 130 fathoms; bottom, volcanic sand. Dredged.

Hemipenæus tomentosus (1 ♀).

New Hebrides. Surface.

Penæus gracilis.
Sergestes rinki.
Sergestes ancylops.

Lucifer reynaudii.
Oodeopus serratus.
Oodeopus armatus.

Oodeopus duplex.

Between Api and Cape York. Surface.

Rhomaleocaris hamulus.

Hectarthropus compressus (1).

STATION 181. August 25, 1874; lat. 13° 15' S., long. 151° 49' E.; depth, 2440 fathoms; bottom, red clay; bottom temperature, 35°·8. Trawled.

Benthesicymus brasiliensis (1 ♂).

AcanthePHYRA sica (1).