

STATION 214. February 10, 1875; lat. 4° 33' N., long. 127° 6' E.; south of the Philippines; depth, 500 fathoms; bottom, blue mud; bottom temperature 41°·8. Trawled.

<i>Pentacheles lævis</i> (1 ♀).		<i>Acanthephyra brachytelsonis</i> (3).
<i>Benthesicymus altus</i> (3 ♂).		<i>Acanthephyra acutifrons</i> (1).
<i>Heterocarpus alphonsi</i> (14 ♀ ♂).		<i>Nematocarcinus undulatipes</i> (3 ♂ ♀).
		<i>Nematocarcinus tenuirostris</i> (5).

STATION 215. February 12, 1875; lat. 4° 19' N., long. 130° 15' E.; near the Philippine Islands; depth, 2550 fathoms; bottom, red clay; bottom temperature, 35°·4. Trawled.

Hymenodora glauca (1).

STATION 218. March 1, 1875; lat. 2° 33' S., long. 144° 4' E.; north of New Guinea; depth, 1070 fathoms; bottom, blue mud; bottom temperature, 36°·4. Trawled.

<i>Paraxius altus</i> (1).		<i>Glyphocrangon granulosis</i> (2 ♀ ♂).
<i>Polycheles helleri</i> (1).		<i>Nematocarcinus tenuipes</i> (1 ♂).
<i>Pentacheles obscura</i> (1).		<i>Nematocarcinus intermedius</i> (2 ♀ ♂).

STATION 219. March 10, 1875; lat. 1° 54' 0" S., long. 146° 39' 40" E.; north of New Guinea; depth, 150 fathoms; bottom, coral mud. Trawled.

<i>Panulirus angulatus</i> (1).		<i>Plesionika spinipes</i> (8 ♂ ♀).
<i>Penæus philippinensis</i> (10 ♂ ♀).		<i>Plesionika unidens</i> (4 ♂ ♀).
<i>Sicyonia lævis</i> (1).		<i>Stylodactylus bimaxillaris</i> (1 ♀).

STATION 220. March 11, 1875; lat. 0° 42' S., long. 147° 0' E.; north of New Guinea; depth, 1100 fathoms; bottom, Globigerina ooze; bottom temperature, 36°·2. Trawled.

<i>Gennadas parvus</i> (3 ♀).		<i>Oplophorus typus</i> (1 ♀).
-------------------------------	--	--------------------------------

North of New Guinea. Surface.

<i>Sergestes corniculum</i> .		<i>Sciacaris telsonis</i> .
<i>Sergestes læviventralis</i> .		<i>Amphion zoea</i> .

STATION 227. March 27, 1875; lat. 17° 29' N., long. 141° 21' E.; surface temperature, 79°·2. Surface.

Eretmocaris longicaulis (1).