

STATION 235. June 4, 1875; lat.  $34^{\circ} 7'$  N., long.  $138^{\circ} 0'$  E.; south of Japan; depth, 565 fathoms; bottom, green mud; bottom temperature,  $38^{\circ}\cdot 1$ . Trawled.

<i>Penaeus tenellus.</i>	<i>Acanthephyra sica</i> (2 ♀ ♂).
<i>Benthesicymus altus</i> (4 ♀ ♂).	<i>Nematocarcinus tenuipes</i> (6 ♂ ♀).
<i>Gennadas parvus</i> (?).	<i>Notostomus japonicus</i> (1 ♂).

STATION 236. June 5, 1875; lat.  $34^{\circ} 58'$  N., long.  $139^{\circ} 29'$  E.; south of Japan; depth, 775 fathoms; bottom, green mud; bottom temperature,  $37^{\circ}\cdot 6$ . Trawled.

<i>Sergestes prehensilis</i> (1 ♂).	<i>Pasiphæa amplidens</i> (1).
<i>Acanthephyra brachytelsonis</i> (1 ♀).	<i>Pasiphæa acutifrons</i> (1).

STATION 237. June 17, 1875; lat.  $34^{\circ} 37'$  N., long.  $140^{\circ} 32'$  E.; near Yokohama; depth, 1875 fathoms; bottom, blue mud; bottom temperature,  $35^{\circ}\cdot 3$ . Trawled.

<i>Aristeus armatus</i> (1 ♀).	<i>Nematocarcinus productus</i> (1 ♀).
<i>Hepomadus glacialis</i> (1 ♀).	<i>Nematocarcinus parvidentatus</i>
<i>Gennadas parvus</i> (1 ♀).	(9 ♂ ♀).
<i>Glypocrangon rimapes</i> (1 ♀).	<i>Nematocarcinus longirostris</i>
<i>Nematocarcinus proximatus</i> (1 ♀).	(10 ♀ ♂).

Off Yokohama.

<i>Penaeus velutinus</i> (6 ♂ ♀).	? <i>Thalassocaris stimpsoni</i> (1). Surface.
-----------------------------------	--

North-west Pacific.

<i>Sergestes corniculum.</i>	<i>Eretmocaris rimapes.</i>
------------------------------	-----------------------------

West Pacific.

<i>Sergestes semiarmis.</i>	<i>Lucifer typus.</i>
-----------------------------	-----------------------

North Pacific.

<i>Sergestes edwardsii.</i>	<i>Lucifer typus.</i>
<i>Amphion reynaudii.</i>	

STATION 245. June 30, 1875; lat.  $36^{\circ} 23'$  N., long.  $174^{\circ} 31'$  E.; depth, 2775 fathoms; bottom, red clay; bottom temperature,  $34^{\circ}\cdot 9$ .

*Hymenodora rostrata* (1). In tow-net at 1700 fathoms.