

BATHYMETRICAL DISTRIBUTION.

TABLE SHOWING THE DEPTH IN FATHOMS AT WHICH THE SPECIES OF EACH GENUS WERE OBTAINED.

	0 to 50.	50 to 100.	100 to 500.	500 to 1000.	1000 to 2000.	2000 to 3000.	3000 to 4000.
Trichobranchiata.							
Aberrantia.							
Pylochelidæ.							
<i>Cheiroplatea</i> ,	1				
Thalassinidæ.							
<i>Thalassina</i> ,	1						
Callianassidæ.							
<i>Callianassa</i> ,	1				
<i>Cheramus</i> ,	1	.	1				
<i>Scallasis</i> ,	1						
Axiidæ.							
<i>Paraxius</i> ,	1		
<i>Eiconaxius</i> ,	1	2			
Thaumastocheleidæ.							
<i>Thaumastocheles</i> ,	1				
Normalia.							
Synaxidea.							
Scyllaridæ.							
<i>Ibaccus</i> ,	1	.	3				
<i>Thenus</i> ,	1						
<i>Arctus</i> ,	1	2	1				
Palinuridæ.							
<i>Panulirus</i> ,	1	.	1				
<i>Palinosyrtus</i> ,	1	.	1				
<i>Phyllosoma</i> ,	3						
Eryonidæ.							
<i>Eryoneicus</i> ,	1		
<i>Polycheles</i> ,	2	1	1		
<i>Pentacheles</i> ,	1	3	2		
<i>Stereomastis</i> ,	1	1			
<i>Willemasia</i> ,	1	1	
Homaridæ.							
<i>Phoberus</i> ,	1			
<i>Nephropsis</i> ,	2			
<i>Nephrops</i> ,	1	1				
Astacidæ.							
<i>Astacopsis</i> ,	{	3 (fresh-water)					