

TABLE VIII.—continued.

HYALONEMATIDÆ.		UNCINATARIA.		INERMIA.		DICTYONINA.	
Depth in Fathoms.	Station.	Depth in Fathoms.	Station.	Depth in Fathoms.	Station.	Depth in Fathoms.	Station.
<i>Semperella schultzei</i> , . . . . .	95	209					
<i>Pheronema globosum</i> , . . . . .	129-140	192	<i>Eureta schmidtii</i> , . . . . .	102	201		
<i>giganteum</i> , . . . . .	129-140	192	<i>semperi</i> , . . . . .	129-140	192		
<i>Semperella schultzei</i> , . . . . .	129-140	192	<i>carleri</i> , . . . . .	129-140	192		
			<i>farreopsis</i> , . . . . .	129-140	192	<i>Myliusia callocyathus</i> , . . . . .	129 140 192
			<i>Hexactinella lata</i> , . . . . .	129-140	192	<i>Aulocystis nitelii</i> , . . . . .	129-140 192
			<i>Cyrtaulon solutus</i> , . . . . .	129-140	192		<i>Dictyonina</i> , . . . . .
			<i>Fieldingia lageloides</i> , . . . . .	129-140	192		129-140 192
<i>Hyalonema (Stylocalgz) apertum</i> , . . . . .	345	232	<i>Farrea clavigera</i> , . . . . .	360	194A	<i>Myliusia callocyathus</i> , . . . . .	{ 360 194A
<i>globus</i> , . . . . .	360	194	<i>Sclerothamnus clausii</i> , . . . . .	360	194A	<i>Aulocystis nitelii</i> , . . . . .	{ 390 24
<i>tozeres</i> , . . . . .	390	24	<i>Aphrocallistes ramosus</i> , . . . . .	375	19		{ 360 194A
			<i>Farrea occa</i> , . . . . .	390	24		
			<i>Aphrocallistes bocagei</i> , . . . . .	390	24		
			<i>Aphrocallistes bocagei</i> , . . . . .	{ 420 344			<i>Dictyonina</i> , . . . . .
				{ 425 343			425
				{ 450 23			207
			<i>Farrea sp.</i> , . . . . .	450	23		
			<i>Chonelasma sp.</i> , . . . . .	450	23		
			<i>Lefroyella decora</i> , . . . . .	435	33		
<i>Hyalonema lusitanicum</i> , . . . . .	600	IV.	<i>Chonelasma lamella</i> , . . . . .	550	148A		
			<i>Hexactinella sp.</i> , . . . . .	550	148A		