TABLE VIII.—*continued.*

Depth in Fathoms.	No. of Dredgings or Trawlings. Total.	With Hexactinellida.	Percentage.	EUPLECTELLIDÆ.	Depth in Fathoms.	Station.	ASCONEMATIDÆ.	Depth in Fathoms.	Station.	ROSSELLIDÆ.	Depth in Fathoms.	Station.
601 to 700	9	4	44·4	*Tægeria pulchra*, . .	610	174c						
				Walteria flemmingii, . .	630	176A				*Euryplegma auriculare*, . .	630	170A
701 to 800	3	...	...									
801 to 900	3	...	...									
901 to 1000	4	2	50									
1001 to 1500	31	7	22·6	*Euplectella nodosa*, . . .	1075	56	*Sympagella nux*, ...	1150	94	*Aulocalyx irregularis*, . .	1075	56
				Rhabdodictyum delicatum, . .	1075	56						
				Euplectella suberea, . . .	1090	V.						
				Holascus fibulatus, . .	1375	146						
				Malacosaccus vastus, . . .	1375	146						
1501 to 2000	35	7	20	*Euplectella suberea*, . . .	1600	124	*Caulophacus latus*, ...	1600	147	*Bathydorus spinosus*, . .	1600	147
				Holascus fibulatus, . .	1600	147				*Aulocalyx irregularis*, . .	1600	147
							Balanites pipetta, . . .	1950	156			
				polejaëvii, . .	1950	157	*Polyrhabdus oviformis*, . .	1975	157			
2001 to 2500	38	8	21·1	*Holascus ridleyi*, . .	2225	211	*Caulophacus latus*, . . .	2050	246	*Caulocalyx tener*, . . .	2025	333
				Dictyocalyx gracilis, . .	2385	261	*elegans*, . . .	2300	241	*Bathydorus fimbriatus*, . .	2300	241
				Malacosaccus unguiculatus, . .	2450	348						
2501 to 3000	33	6	18·2	*Hyalostylus dives*, . . .	2550	289	*Trachycaulus gurlitti*, . .	2550	292			
				Holascus fibulatus, . .	2600	160						
				stellatus, . .	2650	325						
				Euplectella crassistellata, . .	2750	274				*Bathydorus fimbriatus*, . .	2900	248
3001 to 4000	5	...	...									

Zoology Part LIII: Report on the HEXACTINELLIDA. By Professor F. E. Schulze. Bound in Volume 21,1887.

http://19thcenturyscience.org/HMSC/HMSC-Reports/Zool-53/README.htm