

TABLE OF SPECIES OBTAINED BY THE CHALLENGER, SHOWING THEIR GEOGRAPHICAL AND BATHYMETRICAL RANGE.¹

	North Atlantic (Stations 1.-110, and 348-354).	South Atlantic (Stations 111-149, and 317-347).	Southern Ocean* (Stations 144-160).	Indo-Australian (Stations 161-165, and 181-220).	North Pacific (Stations 221-270).	South Pacific (Stations 165-180, and 271-303).	Patagonian (Stations 304-316).	0-50 Fathoms.	50-200 Fathoms.	200-1000 Fathoms.	1000-3000 Fathoms.
<i>Halichondria panicea</i> ,			x		x			x	x		
„ <i>caduca</i> ,							x				
„ <i>solida</i> , n. sp.,						x		30-70			
„ „ var. <i>rugosa</i> , nov.,						x			x		
„ <i>pelliculata</i> , n. sp.,				x					x		
„ <i>latrunculioides</i> , n. sp.,		x								x	
„ sp. (a),		x								x	
„ sp. (b),			x						x		
„ (?) sp. (c),				x				x			
<i>Petrosia similis</i> , n. sp.,		x	x						x		
„ „ var. <i>massa</i> , nov.,							x		x		
„ „ var. <i>compacta</i> , nov.,				x				x			
„ <i>truncata</i> , n. sp.,				x				x			
„ <i>variabilis</i> , var.,				x				x			
„ <i>hispida</i> , n. sp.,			x					x			
<i>Reniera cinerea</i> ,				x					x		
„ <i>implexa</i> , var.,	x									x	
„ <i>aquæductus</i> , var. <i>infundibularis</i> , nov.,							x		x		
„ <i>subglobosa</i> , n. sp.,							x		x		
„ <i>tufa</i> , n. sp.,	x								x		
<i>Pachychalina fragilis</i> , n. sp.,				x				x			
„ <i>melior</i> , n. sp.,				x				x			
„ <i>fibrosa</i> , n. sp.,		?		x				x			
„ <i>lobata</i> ,				x				x			
„ <i>megalorrhaphis</i> , n. sp.,				x					x		
„ <i>elongata</i> , n. sp.,				x				x			
„ (?) <i>punctata</i> , n. sp.,				x				x			
„ (?) <i>pedunculata</i> , n. sp.,			x					10-100			
<i>Chalina palmata</i> ,				x				x			
„ <i>pergamentacea</i> ,				x				x			
„ <i>rectangularis</i> , n. sp.,				x				x			
„ sp. (a),				x				x			
„ sp. (b),				x				x			

¹ This table comprises information derived from the Challenger material only; for fuller information as to the distribution of previously known forms the reader is referred to the Description of Genera and Species.