

PLATE XL.

	PAGE
Fig. 1. <i>Axinella erecta</i> (Station 148A), styli ; $\times 44$,	182
Fig. 1a. Crooked strongyla ("vermicular" spicules) ; $\times 44$	
Fig. 2. <i>Axinella mariana</i> ; a, a, large subtylostylus and stylus ; b, small styli ; $\times 44$,	180
Fig. 3. <i>Axinella profunda</i> , subtylostyli ; $\times 66$,	181
Fig. 3a. Bases of subtylostyli ; $\times 284$	
Fig. 4. <i>Thrinacophora cervicornis</i> ; a, stylus, from one of the surface brushes ; b, oxeote spicule ; $\times 190$,	194
Fig. 4a. Trichodragmata ; $\times 190$	
Fig. 5. <i>Hymeniacidon</i> (?) <i>subacerata</i> , styli ; $\times 66$,	169
Fig. 5a. Base and apex of stylus ; $\times 190$	
Fig. 6. <i>Phakellia flabellata</i> , large styli ; $\times 190$,	171
Fig. 6a. Small styli from surface tuft ; $\times 190$	
Fig. 7. <i>Dendropsis bidentifera</i> ; a, styli ; b, strongylote spicule ; c, stylus with spined base ; $\times 66$,	192
Fig. 7a. Two basally spined styli ; $\times 190$	
Fig. 7b. Bases of three of the basally spined styli ; $\times 284$	
Fig. 7c. Spined oxea ; $\times 284$	
Fig. 8. <i>Raspailia tenuis</i> , two of the large styli ; $\times 66$,	189
Fig. 8a. a, styli ; b, oxeote spicule ; $\times 190$	
Fig. 8b. Strongyla ; $\times 190$	
Fig. 9. <i>Ciocalypta amorphosa</i> ; a, oxea ; b, styli ; $\times 66$,	176