

PLATE XLIII.

| | PAGE |
|---|------|
| Fig. 1. <i>Trichostemma sarsii</i> (Station 73), seen from above; $\times 2$, | 218 |
| Fig. 1a. Same specimen, side view; $\times 2$ | |
| Fig. 2. <i>Trichostemma sarsii</i> , another specimen (Station 73), seen from above; $\times 2$, | 218 |
| Fig. 2a. Same specimen, side view; $\times 2$ | |
| Fig. 3. <i>Trichostemma sarsii</i> (Station 184), seen from above; $\times 2$, | 218 |
| Fig. 3a. Same specimen, side view; $\times 2$ | |
| Fig. 4. <i>Trichostemma irregularis</i> , seen from above; $\times 2$, | 220 |
| Fig. 4a. Same specimen, side view; $\times 2$ | |
| Fig. 5. <i>Trichostemma irregularis</i> , another specimen, side view; $\times 2$, | 220 |
| Fig. 6. <i>Stylocordyla stipitata</i> (Station 49); $\times 2$, | 223 |
| Figs. 7, 8, 9. <i>Stylocordyla stipitata</i> (Station 147); $\times 2$, | 223 |
| Fig. 10. <i>Stylocordyla stipitata</i> , var. <i>globosa</i> (Kerguelen); natural size, | 224 |
| Figs. 11, 12. <i>Suberites elongatus</i> ; $\times 2$, | 205 |
| Figs. 13, 14. <i>Suberites ramulosus</i> , var. <i>cylindrifera</i> ; natural size, | 208 |