

PLATE XLVI.

	PAGE
Fig. 1. <i>Pachychalina</i> (?) <i>punctata</i> , dermal skeleton reticulation surrounding <i>p</i> , one of the pores (?); <i>pr</i> , projection into the pore,	24
Fig. 2. <i>Pachychalina</i> (?) <i>punctata</i> , skeleton arrangement, as seen in vertical section,	24
Fig. 3. <i>Siphonochalina intermedia</i> , skeleton arrangement, as seen in vertical section; <i>a</i> , primary fibres; <i>b</i> , secondary fibres,	30
Fig. 4. <i>Pachychalina lobata</i> , surface section, showing the dermal skeleton reticulation and the pores; <i>d.s.</i> , dermal skeleton; <i>d.m.</i> , dermal membrane; <i>p</i> , pores,	22
Fig. 5. <i>Halichondria latrunculioides</i> , dermal skeleton,	6
Fig. 6. <i>Chalina rectangularis</i> , skeleton arrangement, as seen in vertical section,	28
Fig. 7. <i>Esperiopsis symmetrica</i> , skeleton arrangement, as seen in longitudinal vertical section; <i>a-a</i> , imaginary long axis of sponge; <i>b</i> , a band of spiculo-fibre of the inner series; <i>c</i> , a band of spiculo-fibre of the outer series; <i>d</i> , radiating fibres (part only shown); <i>e</i> , embryos,	77
Fig. 8. <i>Dendropsis bidentifera</i> , skeleton arrangement, as seen in transverse section; <i>a</i> , central axis; <i>b</i> , tufts of small spicules surrounding the large projecting spicules; <i>c</i> , small, spined oxea,	192
Fig. 9. <i>Rhaphidophlus filifer</i> , skeleton arrangement, as seen in vertical section; <i>d.c.</i> , the characteristic dermal crust of spicules,	152