

PLATE XLVII.

	PAGE
Fig. 1. <i>Gelliodes licheniformis</i> , surface section, as seen in alcohol, showing the horny fibre ( <i>h.f.</i> ), . . . . .	48
Fig. 1a. Surface section, as seen in Canada balsam, showing the spicules ( <i>s</i> ), while the horny fibre ( <i>h.f.</i> ) is barely visible.	
Fig. 2. <i>Amphilectus ceratosus</i> , vertical section, showing the horny fibre ( <i>h.f.</i> ), and the arrangement of the spicules, . . . . .	125
Fig. 3. <i>Myxilla rosacea</i> , var. <i>japonica</i> , vertical section, showing the arrangement of the skeleton; <i>d.s.</i> , dermal brushes of spicules; <i>m.s.</i> , main skeleton,	130
Fig. 4. <i>Plumohalichondria mammillata</i> , surface section, showing the arrangement of the dermal skeleton, . . . . .	156
Fig. 4a. Vertical section, showing the plumose skeleton fibres ( <i>p.c.</i> ).	
Fig. 5. <i>Clathria lendenfeldi</i> , section parallel with and near the surface, showing the skeleton arrangement; <i>h.f.</i> , the stout horny fibre echinated by spined styli, . . . . .	148
Fig. 6. <i>Desmacidon</i> (?) <i>ramosa</i> , surface section, showing the radiating tufts of oxea,	107
Fig. 7. <i>Phelloderma radiatum</i> , vertical section, showing the arrangement of the skeleton; <i>c</i> , cortex; <i>s.f.</i> , skeleton fibres; <i>f.b.</i> , foreign bodies; <i>i</i> , isochelæ, . . . . .	114