

STATION 57. Lat. $60^{\circ} 14' N.$, long. $6^{\circ} 17' W.$; 632 fathoms; bottom temperature, $30^{\circ} \cdot 5 F.$

Antedon eschrichti. Also the "Valorous," 1875, Station 1; the "Alert," 1875, Franklin-Pierce Bay; the Challenger, Station 48; and the "Triton," 1882 Station 4.

STATION 74. Lat. $60^{\circ} 39' N.$, long. $3^{\circ} 9' W.$; 203 fathoms; bottom temperature, $47^{\circ} \cdot 6 F.$

Antedon tenella. Also at Stations 51, 54, 55; and 1870, Station 17A; also the "Triton," 1882, Stations 2, 5.

STATIONS not recorded.

Antedon rosacea. Also off the coast of Tunis.

Antedon hystrix. Also the "Triton," 1882, Station 4.

H.M.S. "PORCUPINE," 1870.

STATION 13. Off Cape Mondego; lat. $40^{\circ} 16' N.$, long. $9^{\circ} 37' W.$; 220 fathoms; bottom temperature, $52^{\circ} F.$

Antedon phalangium. Also in the Minch; off Cape Sagres; the Seine Bank, and the Mediterranean.

STATION 17A. Lat. $39^{\circ} 39' N.$, long. $9^{\circ} 39' W.$; 740 fathoms; bottom temperature, $49^{\circ} \cdot 3 F.$

Antedon lusitanica.

Antedon tenella. Also 1869, Stations 51, 54, 55, 74; also the "Triton," 1882, Stations 2, 5.

STATION 25. July 27, 1870; near Cape St. Vincent; lat. $37^{\circ} 11' N.$, long. $9^{\circ} 7' W.$; 374 fathoms; rock; bottom temperature, $53^{\circ} \cdot 5 F.$

Actinometra pulchella. Also at Station 31, and by the Challenger at St. Paul's Rocks; perhaps also at Station 192. Abundant in the Caribbean Sea.

STATION 31. Lat. $35^{\circ} 56' N.$, long. $7^{\circ} 6' W.$; 477 fathoms; clay; bottom temperature, $50^{\circ} \cdot 5 F.$

Actinometra pulchella. Also at Station 25, and at St. Paul's Rocks.