

Bahia ; 7 to 20 fathoms.

Antedon carinata. Also in the Pacific, Eastern Archipelago, and Indian Ocean.

Antedon diibeni.

Actinometra lineata.

Actinometra meridionalis. Abundant in the Caribbean Sea.

STATION 135E. October 18, 1873 ; near Tristan da Cunha ; lat. $37^{\circ} 21' 0''$ S., long. $12^{\circ} 22' 30''$ W.; 1000 fathoms ; hard ground, shells, gravel.

Three Pentacrinoid larvæ.

STATION 135G. October 18, 1873 ; off Tristan da Cunha ; lat. $37^{\circ} 10' 50''$ S., long. $12^{\circ} 18' 30''$ W.; 550 fathoms ; hard ground.

Antedon multispina. Also at Station 344.

Simon's Bay, Cape of Good Hope.

Actinometra parvicirra. Also at Stations 174, 186, Banda, Ternate, Admiralty Islands, and Samboangan.

Off Marion Island ; 50 to 75 fathoms.

Antedon exigua. Also at Station 145.

STATION 145. December 27, 1873 ; off Marion Island ; lat. $46^{\circ} 43' 0''$ S., long. $38^{\circ} 4' 30''$ E.; 140 fathoms ; volcanic sand.

Antedon exigua. Also in 50 to 75 fathoms.

Antedon hirsuta.

STATION 147. December 30 ; between Marion Island and the Crozets, 1873 ; lat. $46^{\circ} 16'$ S., long. $48^{\circ} 27'$ E.; 1600 fathoms ; Diatom ooze ; bottom temperature, $34^{\circ} \cdot 2$ F.

Antedon abyssorum.

Antedon bispinosa.

Antedon remota.

Promachocrinus abyssorum. Also at Station 158.

Bathycrinus aldrichianus.

Hyocrinus bethellianus.