

<i>Antedon prolixa</i> , II.-VII.	<i>Actinometra blakei</i> , IV., V.
<i>quadrata</i> , II.-VI.	<i>discoidea</i> , IV.
<i>rosacea</i> , I., II.	<i>meridionalis</i> , I.-V.
<i>spinifera</i> , IV., V.	<i>pulchella</i> , IV.-VII.
<i>tenella</i> , IV.-VII.	<i>Promachocrinus kerguelensis</i> , II.-IV.
Totals.— <i>Antedon</i> , 15 species ; <i>Actinometra</i> , 4 species.	

TABLE IV.—Species found at depths between 100 and 200 fathoms.

<i>Antedon angustiradia</i> .	<i>Antedon patula</i> .
<i>australis</i> .	<i>phalangium</i> , II.-V.
<i>barentsi</i> .	<i>pourtalèsi</i> , V.
<i>carinata</i> , I.-V.	<i>prolixa</i> , II.-VII.
<i>compressa</i> , III.	<i>pusilla</i> .
<i>defecta</i> , III.-V.	<i>quadrata</i> , II.-VI.
<i>discoidea</i> .	<i>quinquecostata</i> .
<i>duplex</i> , III.-V.	<i>robusta</i> .
<i>eschrichti</i> , I.-VII.	<i>rhomboidea</i> .
<i>exigua</i> .	<i>spinifera</i> , III.-V.
<i>flexilis</i> .	<i>tenella</i> , III.-VII.
<i>granulifera</i> .	<i>tenuicirra</i> .
<i>hageni</i> , III.-V.	<i>Actinometra blakei</i> , III.-V.
<i>hirsuta</i> .	<i>discoidea</i> , III.
<i>longicirra</i> .	<i>meridionalis</i> , I.-V.
<i>manca</i> .	<i>pulchella</i> , III.-VII.
<i>parvipinna</i> .	<i>Promachocrinus kerguelensis</i> , II., III.
Totals.— <i>Antedon</i> , 29 species ; <i>Actinometra</i> , 4 species.	

TABLE V.—Species found at depths between 200 and 350 fathoms.

<i>Antedon brevipinna</i> .	<i>Antedon latipinna</i> .
<i>carinata</i> , I.-IV.	<i>occulta</i> . ²
<i>defecta</i> , III., IV.	<i>phalangium</i> , II.-IV.
<i>duplex</i> , III., IV.	<i>pourtalèsi</i> , IV.
<i>eschrichti</i> , I.-VII.	<i>quadrata</i> , II.-VI.
<i>hageni</i> , III., IV.	<i>similis</i> . ²
<i>inæqualis</i> ¹ ?	<i>spinifera</i> , III., IV.
<i>incisa</i> ¹ ?	<i>tenella</i> , III.-VII.

¹ Most probably obtained from 610 fathoms at Station 174.² Most probably obtained from 210 or 255 fathoms at Station 174.