

<i>Antedon tuberculata</i> . ¹	<i>Actinometra pulchella</i> , III.–VII.
<i>Actinometra blakei</i> , III., IV.	<i>stelligera</i> , ¹ I.
<i>meridionalis</i> , I.–IV.	<i>typica</i> , ¹ I.
<i>parvicirra</i> , ¹ I.	<i>Atelecrinus balanoides</i> , VI.
Totals.— <i>Antedon</i> , 15 species ; <i>Actinometra</i> , 6 species.	

TABLE VI.—Species found at depths between 350 and 500 fathoms.

<i>Antedon acæla</i> .	<i>Antedon porrecta</i> .
<i>aculeata</i> .	<i>prolixa</i> , II.–VII.
<i>angusticalyx</i> .	<i>quadrata</i> , II.–IV.
<i>columnaris</i> .	<i>tenella</i> , III.–VII.
<i>cubensis</i> .	<i>tuberosa</i> .
<i>distincta</i> .	<i>valida</i> .
<i>eschrichti</i> , I.–VII.	<i>Actinometra pulchella</i> , III.–VII.
<i>gracilis</i> .	<i>Atelecrinus balanoides</i> , V.
<i>hystrix</i> .	<i>cubensis</i> .
<i>lævis</i> .	<i>Eudiocrinus atlanticus</i> .
<i>multispina</i> , VII.	<i>Promachocrinus naresi</i> .
Totals.— <i>Antedon</i> , 17 species ; <i>Actinometra</i> , 1 species.	

TABLE VII.—Species found at depths between 500 and 800 fathoms.

<i>Antedon alternata</i> , VIII.	<i>Antedon occulta</i> ² ?
<i>angustipinna</i> .	<i>prolixa</i> , II.–VI.
<i>basicurva</i> .	<i>similis</i> ² ?
<i>breviradia</i> , IX.	<i>tenella</i> , III.–VI.
<i>echinata</i> .	<i>tuberculata</i> ² ?
<i>eschrichti</i> , I.–VI.	<i>Actinometra pulchella</i> , III.–VI.
<i>inæqualis</i> .	<i>parvicirra</i> ² ?
<i>incerta</i> .	<i>stelligera</i> ² ?
<i>incisa</i> .	<i>typica</i> ² ?
<i>lineata</i> .	<i>Atelecrinus wyvillii</i> .
<i>longipinna</i> .	<i>Eudiocrinus japonicus</i> .
<i>lusitanica</i> .	<i>semperi</i> , VIII.
<i>multispina</i> , VI.	
Totals.— <i>Antedon</i> , 15 species ; <i>Actinometra</i> , 1 species.	

¹ Most probably obtained from 210 or 255 fathoms at Station 174.² Only if obtained from 630 fathoms at Station 174, which is improbable.