

*Spicules*.—I. Megascleres. 1. *Strongyle*, 0·8 by 0·016 mm. 2. *Orthotriæne*, rare, 0·48 and over by 0·016 mm., cladi 0·32 mm. long and over.

II. Microscleres. 3. *Sterraster*, spherical, depressed; 0·10 mm. in diameter. 4. *Oxyaster*, with but few slender actines, which are sometimes not more than two in number, and thus indistinguishable from centrotylote microxeas; a single actine of a triradiate form 0·039 mm. in length, a microxea 0·079 mm. in length. 5. *Spherules*, 0·0039 mm. in diameter.

*Colour*.—Blackish-brown above where exposed to the light, whitish below.

*Habitat*.—Sebenico, Adriatic; depth, 7 fathoms.

### Genus 3. *Pachymatisma*, Bowerbank.

The oscules are the single openings of cloacal chones, which lead each into a large cloacal cavity beneath the chonal sphincter. The incurrent chones are cribriporal. The somal microsclere is a microstrongyle.

#### *Pachymatisma johnstonia*, Bowerbank.

*Alcyonium primum*, Diosc. (?), Donati, Hist. Mer. Adriat., 1758.

*Halichondria johnstonia*, Bowerbank, Trans. Micr. Soc. Lond., vol. i. p. 63, pl. vi., 1841.

" " Johnston, Hist. Brit. Spong., p. 198, 1842.

*Pachymatisma johnstonia*, Bowerbank, Mon. Brit. Spong., vol. i. pl. xxvii. fig. 353, 1864; vol. ii. pp. 3, 51, 1866; vol. iii. p. 17, pl. viii. figs. 1-7, 1874.

" " O. Schmidt, Spong. Adriat. Meeres, Suppl., ii. p. 12, 1866.

" " Carter, Ann. and Mag. Nat. Hist., ser. 4, vol. iv. p. 11, 1869.

*Caminus osculosus*,<sup>1</sup> Grube, Mitth. ü. St. Malo u. Roscoff, p. 132, pl. ii. fig. 3, 1873.

non *Pachymatisma johnstonia*, Hansen, Spong. Norweg. North Atlant. Exped., p. 17, pl. iv. fig. 15, 1885.

*Sponge*.—Massive, raised into undulating ridges, sessile, surface smooth; oscules and pores distributed over different areas, the oscules usually occupying the summits of the ridges. Cortex about 1 mm. in thickness, sterrastral layer thin, ectochroite well developed.

*Spicules*.—I. Megascleres. 1. *Strongyle*, usually cylindrical, straight or curved, sometimes irregularly; 0·63 mm. long (B.), 1·03 by 0·013 mm. (S.), 0·924 by 0·013 mm. (R.). 2. *Orthotriæne*, rhabdome conical, oxeate or strongylate, 0·44 mm. long. (B.), 0·636 by 0·016 mm. (S.), 0·44 by 0·019 (R.); cladi conical or cylindrical, oxeate or strongylate, 0·13 mm. long (S.), 0·238 mm. (R.).

II. Microscleres. 3. *Sterraster*, ellipsoidal, 0·12 mm. long. (B.), 0·097 by 0·0838 mm. (S.), 0·9 by 0·071 mm. (R.). 4. *Oxyaster*, actines not numerous, conical, finely spined,

<sup>1</sup> I add here a reference to the following memoir, which I have been unable to consult:—Koehler, R., Remarques sur le genre *Caminus* et sur une éponge voisine du *Caminus osculosus*, Grube (Paris, G. Masson, 1884).