

TABLE OF SPECIES OBTAINED BY THE CHALLENGER, SHOWING THEIR GEOGRAPHICAL AND BATHYMETRICAL RANGE.

	N. Atlantic.	S. Atlantic.	Southern Ocean.	W. Pacific.	E. Pacific.	0-50.	51-200.	201-1000.	1001-2000.
<i>Tetilla sandalina</i> ,	x						x		
„ <i>leptoderma</i> ,		x						x	
„ <i>pedifera</i> ,				x				x	
„ <i>coronida</i> ,			x				x		
„ <i>grandis</i> ,			x			x	x		
„ „ var <i>alba</i> ,			x				x		
„ <i>merguiensis</i> , Carter,				x		x			
„ sp.,				x				x	
<i>Chrotella simplex</i> ,				x			x		
„ <i>macellata</i> ,				x		x			
<i>Cinachyru barbata</i> ,			x			x	x		
<i>Craniella simillima</i> , Bwk.,				x		x			
„ <i>carteri</i> ,		x							
„ <i>pulchra</i> ,				x		x			
„ <i>schmidtii</i> ,	x							x	
<i>Samus anonymus</i> , Gray,		x							
<i>Thenea delicata</i> ,			x						x
„ <i>wrightii</i> ,		x							x
„ <i>grayi</i> ,				x				x	
„ <i>schmidtii</i> ,	x							x	
„ <i>fenestrata</i> , O. Schmidt,	x	x							x
„ <i>wyvillii</i> ,				x					
<i>Pæcillastra schulzii</i> ,			x				x		
„ <i>crassiuscula</i> ,	x						x		
„ <i>laminaris</i> ,				x					
„ <i>tenuilaminaris</i> ,				x				x	
<i>Sphinctrella cribrifera</i> ,	x						x		
„ <i>gracilis</i> ,	x						x		
„ <i>ornatus</i> ,	x						x		
<i>Characella aspera</i> ,		x						x	
<i>Triptolemus cladosus</i> ,				x			x		
<i>Pachastrella abyssii</i> , G. Schmidt,		x					x		
<i>Calthropella simplex</i> ,	x						x		
<i>Myriastræ subtilis</i> ,				x		x			
„ <i>simplicifurca</i> ,				x		x			
„ <i>clavosa</i> , Ridley,				x		x			
„ „ var. <i>quadrata</i> ,				x		x			
„ <i>toxodonta</i> ,				x		x			
<i>Pilochrota hæeckeli</i> ,				x		x			
„ <i>pachydermata</i> ,				x		x	x		
„ <i>gigas</i> ,	x					x			
„ <i>tenuispicula</i> ,	x					x			
„ <i>crassispicula</i> ,		x				x			
„ <i>purpurea</i> , var. <i>longancora</i> ,				x		x			
„ <i>anancora</i> ,		x				x			
„ <i>moseleyi</i> ,				x		x			
„ <i>lendenfeldi</i> ,				x		x			