

	N. Atlantic.	S. Atlantic.	Southern Ocean.	W. Pacific.	E. Pacific.	0-50.	51-200.	201-1000.	1001-2000.
<i>Astrella vosmaeri</i> ,	x	x	x	x	x
<i>Anthastra pulchra</i> ,	x	x	x	x	x	x
" <i>communis</i> ,	.	.	.	x	x	x	x	x	x
" <i>parvispicula</i> ,	.	.	.	x	x	x	x	x	x
" <i>pyriformis</i> ,	.	.	.	x	x	x	x	x	x
" <i>ridleyi</i> ,	x	x	x	x	x	x
<i>Stellella phrixens</i> ,	x	x	x	x	x	x
<i>Tribrachium schmidtii</i> (Weltner), .	.	x	.	x	x	x	x	x	x
<i>Disyringa dissimilis</i> (Ridley), .	.	.	x	x	x	x	x	x	x
<i>Stryphnus niger</i> , .	.	.	x	x	x	x	x	x	x
<i>Psummastra murrayi</i> , .	.	.	x	x	x	x	x	x	x
<i>Amphiurus huxleyi</i> , .	.	.	x	x	x	x	x	x	x
<i>Erylus formosus</i> , .	.	.	x	x	x	x	x	x	x
<i>Caminus sphæroconia</i> , .	.	.	x	x	x	x	x	x	x
<i>Cydonium hirsutus</i> , .	.	.	x	x	x	x	x	x	x
" <i>magellani</i> ,	.	.	x	x	x	x	x	x	x
" <i>glariosus</i> ,	.	.	x	x	x	x	x	x	x
" <i>eosaster</i> , .	.	.	x	x	x	x	x	x	x
<i>Synops neptuni</i> , .	.	x	x	x	x	x	x	x	x
" <i>nitida</i> , .	.	x	x	x	x	x	x	x	x
" <i>vosmaeri</i> , .	.	x	x	x	x	x	x	x	x
<i>Isops pachydermata</i> , .	x	x	x	x	x	x	x	x	x
<i>Thrombus challengerii</i> , .	.	.	x	x	x	x	x	x	x
<i>Theonella swinhonis</i> , Gray, .	.	.	x	x	x	x	x	x	x
<i>Discodermia discifurca</i> , .	.	x	x	x	x	x	x	x	x
" <i>panoplia</i> ,	.	x	x	x	x	x	x	x	x
" <i>ornata</i> , .	.	x	x	x	x	x	x	x	x
<i>Neosiphonia superstes</i> , .	.	.	x	x	x	x	x	x	x
<i>Corallistes typus</i> , O. Schmidt,	.	x	x	x	x	x	x	x	x
" <i>masoni</i> (Bow.), .	.	x	x	x	x	x	x	x	x
" <i>thomasi</i> , .	.	x	x	x	x	x	x	x	x
" <i>bowerbanki</i> (Johnson), .	.	x	x	x	x	x	x	x	x
<i>Pleroma turbinatum</i> , .	.	.	x	x	x	x	x	x	x
<i>Callipelta ornata</i> , .	.	.	x	x	x	x	x	x	x
<i>Scleritoderma flabelliformis</i> , .	.	.	x	x	x	x	x	x	x
<i>Siphonidium capitatum</i> , .	.	.	x	x	x	x	x	x	x
<i>Azorica pfeifferae</i> , Carter, .	.	x	x	x	x	x	x	x	x
" <i>marginata</i> , .	.	x	x	x	x	x	x	x	x
" <i>crassiuscula</i> , .	.	x	x	x	x	x	x	x	x
<i>Tretolophus paniceus</i> , .	.	x	x	x	x	x	x	x	x
	15	17	6	47	3	35	33	16	5