

1851. LILJEBORG, WILH.

Bidrag till Norra Rysslands och Norriges fauna, samlade under en vetenskaplig resa i dessa länder 1848. Inlemnad d. 11 December 1850. Kongl. Vetenskaps-Akademiens Handlingar för år 1850. Sednare Afdelningen. (This continues apparently Kongl. Vetenskaps-akademiens Handlingar, för år 1849. Stockholm, 1851.) pp. 233–341.

At page 311 he mentions having observed "vid Schuretskaja i Ryska Lappland," among the Crustacea, "*Gammarus locusta*, Mont., Kröy.;" "*Anonyx Edwardsii*, Kröy.;" "*Caprella lobata* (Muell.); Kröy."

At page 346 occurs the heading, "Förteckning öfver de af mig i trakten af Tromsö i Norge observerade Däggdjur, Foglar etc." Among the Tromsö Crustacea are the following Amphipoda :—" *Gammarus locusta*, Mont. ; Kröy. Varietas : Antennæ superiores inferioribus longiores, et earum pedunculi articulum penultimum pedunc. antenn. infer. excedentes.—*Pardalisea cuspidata*, Kröy.—*Pontoporeia femorata*, Kröy.—*Amphithoe albomaculata*, Kröy.—*Anonyx ampulla* (Phipps) ; Kröy.—*Ischyrocerus minutus*, n. sp. Antennæ superiores inferioribus insigniter longiores, flagello ultimum pedunculi articulum æquante, circiter 6-articulato, articulis elongatis, flagello appendiculari vix dimidiata primi articuli flagelli proprii longitudinem æquante; ultimus pedunculi antennarum inferiorum articulus flagellum circ. 5-articulatum æquans; manus primi et secundi pedum paris fere æquales, subangustæ. Forma minima. Tantummodo specimen unum, feminam oviferam, circ. 4 mm. longam e profundo majore accepi.—*Leucothoe norvegica*, n. sp. (Tab. xx. fig. 4)." For the description, see Note on Liljeborg, 1850. *Ischyrocerus minutus* is identified by Boeck with *Podocerus anguipes*, Krøyer, but it is more probably the same as the species described by Sars in 1882 as *Podocerus minutus*, n. s.

After some Isopods, Liljeborg also mentions "*Caprella lobata* (Muell.); Kröy."

1851. LILJEBORG, WILHELM.

Norges Crustacéer. Crustacea, a clariss. M. W. v. Dueben in Norvegia ad Christiansund et Bergen 1843–44 collecta. Öfversigt af Kongl. Vetenskaps-Akademiens Förfhandlingar. Årg. 8. 1851. No. 1. Stockholm. pp. 19–25.

Of the hundred and six species collected by v. Düben, numbers 34 to 70 belong to the Amphipoda. Many are merely named. On a few, notes are given as follows :—

" 34. *Orchestia litorea*, M. E.—Femina a mari tam diversa forma pedum secundi paris, ut illa formam typicam generis *Talitri* M. E. et hic eandem formam gen. *Orchestiae* efficere videatur. Femina *Talitro tripudianti* Kr. sat similis est; forma pedum secundi paris. prorsus eadem est, longitudo vero pedum quarti et quinti paris diversa, quum hi pedes iisdem secundi paris longiores."

" 42. *Anonyx norvegicus* n. sp.—*A. plauto* Kr. sat. affinis, diversus tamen : antennis superioribus feminæ et maris fere æqualibus, flagello circ. 14-articulato, flagello appendiculari 5-articulato, artic. primo ceteris vix longiore; antennis inferioribus superioribus insigniter —maris plus quam duplo—longioribus; epimero quarto postice profunde sinuato, epimero quinto mediocri, subquadrato, antecedentibus humiliore. Præterea inter se similes." This in 1865 he identifies with *Anonyx gulosus*, Krøyer.

" 51. *Amphithoe Pausilipii*? M. E.—Omnino ei similis, oculi vero non visibles."