

*First Peræopods.*—Side-plates similar to the preceding pair. Branchial vesicles expanding from a narrow neck so as to be widest distally, as long as the first joint, and almost as wide as long. First joint extending a little below the side-plate, second joint very short, third longer than fourth or fifth, with some setæ on the hind margin and apex of front; fourth a little longer and much broader than fifth, with setæ on both margins; fifth not broad and not tapering, with setæ on both margins, those on the straight hind margin short; the finger very short, with short cilia near the hinge and near the nail; the nail with a pointed projecting cap.

*Second Peræopods.*—Side-plates not much longer but very much broader than the preceding pair, the excavation behind descending a very small distance. The first joint not reaching the end of the side-plate, the limb in other respects scarcely differing from the preceding pair.

*Third Peræopods.*—Side-plates broad but not deep, the hind lobe less broad than the front, of about the same depth, crenulate and ciliated round the lower part of its hind margin. The first joint large, oblong, oval, rather broader below than above, with small setæ in the crenulation of the hind rim, and longer setæ on the front margin; second joint very small; third of great size, nearly as long as the first joint, much broader than the fourth, somewhat decurrent, with setæ, some of which are spiniform, along both margins; the fourth joint longer and much broader than the fifth, with setæ on both margins; the fifth not so long as the straight, slender finger, with setæ on both margins, and at the front apex close to the hinge of the finger a group of spines, two short and stout, and a third half the length of the finger, of great strength; the finger tapering, minutely pectinate in front, the nail spine-like, with a cilium at its base, and sheltered by a long cap, the peak of which projects beyond it.

*Fourth Peræopods.*—The hind lobe of the side-plates produced much below the front one. The first joint more rounded than in the preceding pair, rather broader, but rather shorter, especially behind; the other joints similar but longer, the third and fourth also wider, the third more strongly armed with spines.

*Fifth Peræopods.*—Side-plates small, not bilobed. Branchial vesicles small, ovate, a little larger than the side-plates. First joint broader and behind much longer than that of the preceding pair; the third joint large, broad, and strongly spined; fourth joint longer than in the preceding pairs, with spines as well as setæ on the front margin; the fifth joint equal in length to the finger, the dorsal cilium of which in this, as in the two preceding pairs, is very small.

*Pleopods.*—Groups of setæ on the peduncles, two hooked spines both apically sharp, one with three, the other with two retroverted teeth, the opposite margins with backward serrature; the rami with fourteen joints to the inner, sixteen to the outer; the first joint of the inner with three cleft spines at the upper part, some plumose setæ below, and groups of setæ on the opposite margin.