

1884. *Moera*, Chilton, Proc. Linn. Soc. N.S.W., vol. ix. pt. iv. (extract), p. 3.
 1884. „ Miers, Report on Zool. Coll. H.M.S. "Alert," p. 315.
 1885. *Mæra*, Carus, Prodrömus Faunæ Mediterraneæ, p. 414.
 1885. *Megamoera*, Haswell, Proc. Linn. Soc. N.S.W., vol. x. pt. i. (extract), p. 9.
 1885. *Moera*, Haswell, Proc. Linn. Soc. N.S.W., vol. x. pt. i. (extract), p. 11.
 1885. *Mæra*, Sars, Den norske Nordhavs-Exp., p. 177.
 1886. „ Gerstaecker, Bronn's Klassen und Ordnungen, Bd. v. Abth. ii. p. 510.
 1887. „ Chevreux, Catalogue des crust. Amph. Bretagne, p. 19.

For the original definition of the genus *Mæra*, see Note on Leach, 1813 (p. 84), for that of *Ceradocus*, see Note on Costa, 1857 (p. 298), for that of *Leptothoe*, see Note on Stimpson, 1854 (p. 277), and for that of *Megamoera*, see Note on Spence Bate, 1862 (p. 335). Boeck's definition of *Mæra* is as follows:—

“Mandibles with the third joint of the palp narrow, not very elongate.

“First Maxillæ with the inner plate narrow and apically furnished with few plumose setæ.

“Body elongate, narrow.

“Side-plates little; the fourth pair scarcely larger than the fifth.

“Upper Antennæ much longer than the lower, the peduncle elongate.

“The legs slender, elongate; the *Second Gnathopods* with the hand much larger than in the first pair; the *Third, Fourth, and Fifth Peræopods* with the first joint more or less dilated.

“The *Third Uropods* extended beyond the *First* and *Second* pairs; the rami elongate, narrow, very setose on the margin; the inner ramus only a little shorter than the outer.

“The *Telson* not very elongate, deeply cleft.”

In *Mæra longimanus*, Leach, the third joint of the mandibular palp is longer than the second, and in *Mæra rubromaculata* (Stimpson) the inner plate of the *First Maxillæ* has the plumose setæ numerous instead of few, and not confined to the apex. The epithet “few” should, I think, be omitted from the generic character of the *First Maxillæ*.

Mæra rubromaculata, Stimpson (Pls. XCV., XCVI.).

- 1855-6. *Gammarus rubro-maculatus*, Stimpson, Proc. Acad. Nat. Sci. Philad., vol. vii.
 1862. *Megamoera serrata*, Sp. Bate, Brit. Mus. Catal. Amph. Crust., p. 227, pl. xxxix. fig. 5.
 1880. *Melita* (?) *Ramsayi*, Haswell, Proc. Linn. Soc. N.S.W., vol. iv. p. 246, pl. x. fig. 1.
 1880. *Moera rubro-maculata* (Stimpson), Haswell, Proc. Linn. Soc. N.S.W., vol. iv. p. 267, pl. x. fig. 4.
 1880. „ *spinosa*, Haswell, Proc. Linn. Soc. N.S.W., vol. iv. p. 268, pl. x. fig. 5.
 1880. „ *Ramsayi*, Haswell, Proc. Linn. Soc. N.S.W., vol. iv. p. 334.
 1882. „ „ Haswell, Catal. Australian Crust., p. 253.
 1882. „ *rubro-maculata*, Haswell, Catal. Australian Crust., p. 254.