

standing unless taken in connection with the definition of the genus "*Thaumatops*," given in the same work as follows:—

"Head very large, bordered with a serrated crest around the middle. First two pairs of pereiopoda [*First and Second Gnathopods*] cheliform. Second pair of uropoda are totally wanting. The interior rami are coalesced in both existing pairs of uropoda."

### Genus *Cystisoma*, Guérin-Méneville, 1842.

- 1842. *Cystisoma*, Guérin-Méneville, Revue zoologique, juillet 1842, p. 214.
- 1852. " Dana, Amer. Journ. Sci. and Arts, ser. 2, vol. xiv. No. 41.
- 1852. " Dana, U.S. Explor. Exped., vol. xiii. pt. ii. pp. 981, 1442.
- 1862. *Cystosoma*, Spence Bate, Brit. Mus. Catal. Amph. Crust., p. 311.
- 1873. *Thaumops*, v. Willemoes Suhm, Proc. Roy. Soc. Lond., vol. xxi. p. 206.
- 1874. " v. Willemoes Suhm, Phil. Trans. Roy. Soc. Lond., vol. clxiii. pp. 629, 637.
- 1874. *Cystosoma*, v. Willemoes Suhm, Nature, vol. ix. p. 182.
- 1875. *Thaumatops*, von Martens, Zool. Record for 1873, vol. x., Crustacea.
- 1875. *Cystosoma*, v. Willemoes Suhm, Zeitschr. f. wiss. Zool., Bd. xxv. p. 37.
- 1875. *Cystisoma*, v. Willemoes Suhm, Trans. Linn. Soc. Lond. (Zool.), ser. 2, vol. i. pt. i. p. 24.
- 1876. " v. Willemoes Suhm, Proc. Roy. Soc. Lond., vol. xxiv. p. 570.
- 1877. *Cystosoma*, Wyville Thomson, The Atlantic, vol. i. p. 129.
- 1879. " Pagenstecher, Sammlung gem. wissenschaftl. Vorträge, ser. xiv. Hft. 315, 316, p. 39.
- 1885. *Cysteosoma*, Bovallius, Mimonectes, A Remarkable Genus of Amph. Hyper., p. 2.
- 1885. *Cystosoma*, Stebbing, Narr. Chall. Exp., part ii. p. 622.
- 1886. *Thaumatops*, Bovallius, On the genus Cysteosoma or Thaumatops, Bihang till K. Svensk. Vetensk. Akad. Handl., Bd. 11, No. 9.
- 1886. *Cystosoma*, Gerstaecker, Bronn's Klassen und Ordnungen, Bd. v. Abth. ii. p. 490.
- 1887. *Thaumatops*, Bovallius, Systematical List of Amph. Hyper., Bihang till K. Svensk. Vetensk. Akad. Handl., Bd. 11, No. 16, p. 14.
- 1887. " . Bovallius, Arctic and Antarctic Hyperids, Vega-Exped., Bd. iv. p. 557.

For the original definition of the genus *Cystisoma*, see Note on Guérin-Méneville, 1842 (p. 196); this may be compared with the account of the earliest described species, *Oniscus spinosus*, in the Note on J. C. Fabricius, 1775 (p. 40). For the definitions successively given for *Thaumops*, see Notes on v. Willemoes Suhm, 1873 (p. 423), 1874 (p. 440). The definition of the genus under the name *Thaumatops* given by Bovallius in 1887 has been quoted above; in 1886 he defined it as follows:—

"The body is hyaline, the segments distinctly separated from one another (except the two first pereional segments of *Th. Neptunus* and *Th. pellucida*, which are coalesced). The epimerals are only indicated.

"The head is large, more or less rounded, tumid. The eyes are large, occupying almost all the upper surface of the head.

"Only the first pair of antennæ are developed, straight, few-articulated. The second pair are represented by two small tubercles.