

*Second Peræopods.*—Side-plates rather broader than the preceding pair, of somewhat pentagonal shape, but with all the angles rounded. The limb nearly as in the preceding pair, but the first joint more regularly oval, the third joint rather longer and the fourth rather shorter.

*Third Peræopods* rather shorter than the two preceding pairs. Side-plates broader than deep, rather deeper in front than behind. The first joint with convex front margin, the hind margin of the front surface almost straight till near the rounded apex, which projects behind the other margin and the following joint; some of the muscles almost reach the base of the joint; the following joints similar to those of the second peræopods, but without spines or pectination, the fourth joint less wide, the fifth joint shorter but fully as broad; the finger similar.

*Fourth Peræopods.*—The side-plates broader than the preceding pair, very much broader than deep. The branchial vesicles longer and broader than the first joint. The limb scarcely differing from the preceding, but the first joint rather broader and more oval, the third and fourth joints subequal in length, the fifth joint a little longer than in the third peræopods.

*Fifth Peræopods* like the fourth, but the side-plates less deep, the first joint rather larger and more oblong, the fifth joint shorter than in the third pair. The third, fourth, and fifth peræopods are as nearly as possible equal to one another, but shorter than the first and second.

*Pleopods.*—The cleft spine has the serrate arm much stouter than the other, which is also shorter and has a very small subapical dilatation; the first joint of the inner ramus, besides being ciliated above the cleft spine, has some five or six plumose setæ below it; the first joint of the outer ramus is also bordered with five or six plumose setæ; each ramus has from eighteen to twenty joints.

*Uropods.*—Peduncles of the first pair longer but narrower than those of the third, longer than the rami, the distal margin minutely pectinate; the rami set a little apart at the base, elongate lanceolate, the adjacent edges pectinate, near the base slightly emarginate and furred, the other edges smooth, the apices acute; the outer ramus a little shorter than the inner; the peduncles of the second pair a little shorter and narrower than those of the third, longer than the rami; the outer ramus shorter and much narrower than the inner, its inner margin pectinate, the inner ramus a little shorter than the rami of the first pair but much broader, the outer margin and lower part of the inner margin pectinate; this ramus scarcely reaches the middle of the rami of the third pair; peduncles of the third pair much longer than the rami, very broad except near the base, the lower part of the inner margin pectinate, its apex sharply squared; the rami subequal in length, the outer much the narrower, with its outer margin smooth, the inner pectinate; the inner ramus broadly lanceolate, with both edges pectinate.

*Telson* longer than broad, not half the length of the peduncles of the third uropods,