

*Second Gnathopods* very like the first, but the first joint longer and rather more slender, and the wrist widening distally, with straight or slightly concave distal margin forming the widest part of the joint and projecting much behind the hand, the hind margin furnished with five setules or spinules successively larger, and three submarginal hairs or setules, besides a little excessively fine furring.

*First Peræopods*.—The first joint narrowest at the neck, the front margin straight; the second joint longer than broad; the third joint subequal in length to the fourth, which is a little narrower and slightly curved; the fifth joint longer and narrower than the fourth, slightly curved, the concave hind margin of this and the preceding joint faintly furred; the finger bulbous at the base, very slightly curved, less than half the length of the fifth joint. There are microscopic setules on various parts of the limb.

*Second Peræopods* like the first, except that they are slightly stouter, and the third, fourth, and fifth joints are longer.

*Third Peræopods* much longer than the preceding pairs, the first joint not widely expanded, more than twice as long as broad, with some small spines along the upper part of the front margin which below is weakly serrate; the second joint longer than broad; the third and two following joints much more elongate than in the preceding pair, the third rather longer than the fourth; the fifth slender, curved, much longer than either of the two preceding joints, nearly as long as the first; the finger small, about a sixth of the length of the preceding joint.

*Fourth Peræopods* shorter than the third, the first joint rather longer and at the upper part much more widely expanded, the front margin produced a little below the hinder, and having a few spines, the convex hind margin smooth; the second and third joints rather larger than in the preceding pair, the third joint having its distal margin finely pectinate and armed on either side with a couple of spines; the fourth joint about half the length of the third, the distal armature similar, but the pectination much stronger, the front margin very minutely pectinate at the middle, but at a little distance from the apex carrying four short broad teeth; the fifth joint longer than the third, curved, the concave front margin strongly pectinate; the finger small.

*Fifth Peræopods*.—The first joint narrowly pear-shaped, not so long as the first joint of the third pair, but as broad at its broadest part; the second joint not longer than broad; the third about twice as long as the second; the fourth nearly twice as long as the third, but narrower; the fifth nearly as long as the fourth; the minute finger projecting in front of the fifth joint and strongly bent, so that the linear or setiform termination is brought very near to the base and projects in advance of it.

*Pleopods*.—Coupling spines minute; cleft spine having the arm with the subapical dilatation considerably longer than the other; the first joint of the inner ramus carrying five setæ below the cleft spine; the first joint of the outer ramus longer than