

## Caprella—continued.

<i>leptonyx</i> , Heller, 1866, . . . . .	367
	18, fig. 7, 24, fig. 9, 30 fig. 11, 68, 72, 73, 85 95, 98, 121, 124, 129, 157 180, 183, 191, 192, 196 203, 205, 206, 219, 220 224, 249, 254, 270, 280 282, 342, 390, 397, 410 421, 457, 479, 480, 511 528, 536, 548, 1621 1635
<i>linearis</i> , Bosc, 1802, . . . . .	478, 479
	72, 147, 163, 203, 217 221, 235, 243, 251, 252 276, 282, 284, 319, 327 480, 528, 1645
<i>liparotensis</i> , Haller, 1879, . . . . .	232, 233, 276
<i>lobata</i> , Latreille, 1802, . . . . .	397
<i>longicollis</i> , Nicolet, 1849, . . . . .	276
<i>longicornis</i> , Boeck, 1870, . . . . .	212, 282
<i>longimanus</i> , Stimpson, 1854, . . . . .	397
<i>longispina</i> , Krøyer, 1845, . . . . .	288
Lovèni, Boeck, 1870, . . . . .	95, 124, 191
<i>luctator</i> , Stimpson, 1855-6, . . . . .	385, 429
<i>mantis</i> , <sup>1</sup> Latreille, 1816, . . . . .	498, 571
<i>mediterranea</i> , <sup>2</sup> Mayer, 1881, . . . . .	1644, 1646
<i>megacephala</i> , A. Milne-Edwards, 1868, . . . . .	330
<i>microtuberculata</i> , G. O. Sars, 1879, . . . . .	367, 390
var. <i>spinigera</i> , Hansen, 1887, . . . . .	247
<i>Modesta</i> , Herklots, 1861, . . . . .	169, 191, 388, 429, 1257
<i>monacantha</i> , Heller, 1866, . . . . .	480
<i>nichtensis</i> , Brandt, 1851, . . . . .	329, 444
<i>nodosa</i> , Templeton, 1836, . . . . .	513
Novæ-Zelandiæ, Kirk, 1878-9, . . . . .	367
<i>obesa</i> , P. J. van Beneden, 1861, . . . . .	265
<i>obesa</i> , Haswell, 1880, . . . . .	85
<i>obtusa</i> , Heller, 1866, . . . . .	119, 157, 254
<i>obtusirostris</i> (see <i>cornuta</i> ), . . . . .	74, 85, 105, 119, 124 157, 183, 191, 206, 223 249, 254, 282, 290, 306 521
Penantis, <sup>3</sup> Leach, 1813-14, . . . . .	375
<i>Pennantii</i> , Fleming, 1823, . . . . .	323, 397
	98, 124, 129, 249
<i>protelloides</i> , Czerniavski, 1868, . . . . .	180
<i>punctata</i> , Boeck, 1860, . . . . .	1244, 1627
<i>punctata</i> , Risso, 1816, . . . . .	265, 349
<i>quadriloba</i> , Krøyer, 1838, . . . . .	276
<i>quadrispinis</i> , Grube, 1864, . . . . .	195, 207, 276
<i>robusta</i> , Dana, 1852, . . . . .	169, 191, 265, 385, 388
<i>robusta</i> , Stimpson, 1854, . . . . .	429, 1257, 1629, 1633
<i>sanguinea</i> , Gould, 1841, . . . . .	pl. cxliv
<i>scaura</i> , Templeton, 1836, . . . . .	66, 105, 140
<i>scolopendroides</i> , Lamarck, 1801, . . . . .	

## Caprella—continued.

<i>scolopendroides</i> , Ross, 1826, . . . . .	180, 468, 1619, 1620
	8, fig. 4, 180, 197, 203 217, 271, 302, 319, 323 358, 367, 397, 430, 450 483, 536, 1634, 1645
<i>septentrionalis</i> , Krøyer, 1838, . . . . .	288
<i>solitaria</i> , Stimpson, 1855-6, . . . . .	281, 358, 468
<i>spinifera</i> , Bell and Westwood, 1855, . . . . .	233
<i>spinifrons</i> , Nicolet, 1849, . . . . .	1644, 1646
<i>spinigera</i> (see <i>microtuberculata</i> ), . . . . .	195, 221, 306
<i>spinosa</i> , Goodsir, 1842, . . . . .	443, 1259, 1632
<i>spinosa</i> , Lockington, 1875, . . . . .	468, 536
<i>spinosissima</i> , Sp. Bate, 1862, . . . . .	276, 277, 422, 459, 536 571, 585, 599
<i>spinosissima</i> , Norman, 1873, . . . . .	254, 536
<i>spinulata</i> , Couch, 1852, . . . . .	276
<i>Stimpsoni</i> , Sp. Bate, 1862, . . . . .	265
<i>subtenuis</i> (see <i>attenuata</i> ), . . . . .	230, 367
<i>tabida</i> , Lucas, 1849, . . . . .	511
<i>tenella</i> , <sup>4</sup> Sp. Bate, 1862, . . . . .	163, 191, 221, 224, 430 1259, 1629
<i>tenuis</i> , Haswell, 1880, . . . . .	195, 196, 221, 224, 282 306, 331, 536, 547
<i>tuberculata</i> , Guérin 1836, . . . . .	344
<i>tuberculata</i> , Goodsir, 1842, . . . . .	343
<i>tuberculosa</i> , Kinahan, 1863, . . . . .	19, 338
<i>typica</i> , Sp. Bate, 1862, . . . . .	66, 68, 73
<i>ultima</i> , Sp. Bate, 1862, . . . . .	410, 1629
<i>ventricosa</i> , Lamarck, 1801, . . . . .	
<i>verrucosa</i> , Boeck, 1872, . . . . .	
<i>Caprellina</i> (now <i>Caprellinopsis</i> )—	
<i>longicollis</i> , Mayer, 1882, . . . . .	233, 499, 535
Mayeri, Pfeffer, 1888, . . . . .	1653
<i>Novæ-Zelandiæ</i> , G. M. Thomson, . . . . .	499, 535
<i>Caprellinoides</i> —	
<i>tristanensis</i> , Stebbing, 1888, . . . . .	1238, pl. cxli
<i>Caprellinopsis</i> (see <i>Caprellina</i> )—	
<i>longicollis</i> , Stebbing, 1888, . . . . .	233, 1237
<i>Capreola</i> —	
<i>Linearis</i> , de Brébisson, 1825, . . . . .	121
<i>Carcinococcus</i> —	
<i>Costæ</i> , Costa, 1851, . . . . .	248, 250, 369
<i>Ovatus</i> , Costa, 1851, . . . . .	248, 250
<i>Poveriæ</i> , Costa, 1851, . . . . .	248, 250
<i>Carcinornis</i> —	
<i>acutirostris</i> , Costa, 1864, . . . . .	347
<i>inflaticeps</i> , Costa, 1864, . . . . .	347
<i>Cardenio</i> —	
<i>paurodactylus</i> , Stebbing, 1888, . . . . .	806, pl. liii
<i>Cephalaspis</i> —	
<i>Seticauda</i> , Costa, 1851, . . . . .	249
<i>Ceradocus</i> —	
<i>orchestipes</i> , Costa, 1853, . . . . .	206, 299, 367, 561
<i>Cerapodina</i> —	
<i>abdita</i> , M.-Edw., 1840, . . . . .	168, 189, 462

<sup>1</sup> On p. 95, lines 29, 30, for "ou en distingue," read "on en distingue," and for "de" read "du."

<sup>2</sup> A catalogue name, Mitthell. Zool. Stat. Neapel 1881, Bd. 2 p. 526. See Mayer, Die Caprelliden, p. 51, where it is recognised as a synonym of *Caprella dentata*, Haller.

<sup>3</sup> As Mayer identifies this with *acutifrons*, Leach-Latreille, 1816, it would seem to take precedence of the latter name.

<sup>4</sup> This is *Ægina ? tenella*, Dana, which Boeck named *Æginella tenella*.